

**Virucidal activity of**  
**„PMF-concentrate“**  
**against the *Transmissible Gastroenteritis Virus of***  
***Swine (TGEV)***  
**(used as a model virus for the**  
***Middle East Respiratory Syndrome Coronavirus (MERS-CoV)***

**Short report of the screening test S3**

by

**PD Dr. Olaf Thraenhart and Dr. Christian Jursch**

**Study time: in March 2015**

**Principal:** PMF Natural Products company  
 Arab Republic of Egypt

**Product:** **PMF-concentrate**  
 [Lot-no.: not specified; product sample as arrived; Arrival: 01.08.2014, Storage at 2-8°C]

**Parameter of test:**

- 0,75 g of PMF-concentrate solved within 2,95 mL of A. bidest (25,42% [w/v])
- T = 37° C and 60 and 240 min. of exposure

**Test system:**

- Transmissible Gastroenteritis Virus of Swine (TGEV); Strain: Toyama  
 (Origin: Virusbank of the Bundesforschungsanstalt f. Viruskrankheiten der Tiere; Friedrich Löffler-Institut, Insel Riems, Germany)
- ST75/2 cells (foetal testis cells of swine)  
 (Origin: Robert Koch-Institute, Berlin, Germany)

**Test method:**

- The testing was performed following the guideline of the DVV and the Robert Koch-Institute (DVV/RKI-guideline [*Bundesgesundhbl.* (2008); 51 (8):937-945]): for the quantitative virucidal suspension test (QST).
- With this testing virus titration of the main samples was performed according to Lycke's methodology (*Arch Ges Virusforsch* (1957); 7:483-493).

**Tab. 1: Dosage of product (solvent: A. bidest)**

Set	Product(s)	Conc. in Test (Ix)	Working sol. (x 1,25)	Dosage	pH of working sol.
#1	PMF-concentrate	20,34%	25,42%	0,75 g in 2,95 mL	pH 9,54 (in test: pH 9,52 )

**Tab. 2: Results of virus inactivation**

Samples	1a + 1b	2a + 2b
	Virus inactivation	
Exposure time	t = 60 min.	t = 240 min.
Virus input <sup>1</sup> (per test volume)	6,11 ± 0,34	5,50 ± 0,32
Detection limit (cytotoxicity level)	< 1 ID <sub>50</sub> (0,0 lg ID <sub>50</sub> )	
Residual virus <sup>1</sup> (log ID <sub>50</sub> ± K [95%])	1,80 ± 0,34 (43/480 virus positive cell culture units)	< 1 ID <sub>50</sub> (0,0 lg ID <sub>50</sub> ) (0/480 virus positive cell culture units)
<b>Reduction <sup>2</sup></b> (log ID <sub>50</sub> ± K [95%])	<b>4,31 ± 0,48</b>	<b>&gt; 5,50 ± 0,45</b>

<sup>1</sup> = Calculation of 95% confidential interval of virus titer as well as virus reduction following DVV/RKI-Guideline.

<sup>2</sup> = Virus reduction: titer of virus control minus titer of sample (lg ID<sub>50</sub>).

**Results: (cf. Tab. 2)**

**Control tests**

- With S3 the amount of input virus at 37° C was estimated to  $\lg ID_{50} = 6,13 \pm 0,34$  after 60 min.
- After 240 min. of incubation at 37° C virus titer was declined to  $\lg ID_{50} = 5,52 \pm 0,32$  due to the influence of temperature ( $\Delta$  virus titer =  $0,61 \pm 0,46$ ).
- With the *Lycke's* method a sample dilution was done (VF = 1000). With that dilution no cytotoxicity was visible and the susceptibility of the detection cells was given ( $\Delta$  titer =  $0,18 \pm 0,47$ ).

**Virus inactivation**

- With 20,34% of PMF (final test concentration) and after 60 min. the virus reduction factor was estimated to **RF = 4,31 ± 0,48**.
- After the exposure time was prolonged to 240 min. no residual virus could be detected. The corresponding virus reduction factor was estimated to **RF > 5,50 ± 0,45**.

**Conclusions:**

- Prolongation of the exposure at 37° C from 60 to 240 min. was associated with only a minor reduction of input virus ( $\Delta$  virus titer =  $0,61 \pm 0,46$ ). The test virus was sufficiently stable at the test temperature over the observation period.
- After 60 min. at 37° C the tested product PMF-concentrate in its 20,34% dilution (final concentration) inactivated *TGEV* by RF =  $4,31 \pm 0,48$  or by 99,995% under the test conditions. After 240 min. a virus reduction factor of RF >  $5,5 \pm 0,45$  was observed, correspondent to a virus reduction of 99,999%.

Luckenwalde, 18<sup>th</sup> of March 2015

Dr. Christian Jursch  
(Laboratory manager and Managing Director of Eurovir)

Inactivation of *TGEV* (a model virus for MERS-CoV) by PMF-concentrate  
- testing with the quantitative virucidal suspension test at T = 37 °C -




Fig. 1

**- Attachment: Experimental protocols -**

- Virucidal activity of the product *PMF-concentrate* - Experiment S3 at T = 37° C / testing in the *quantitative suspension test (QST)* using the methodology of *Lycke* for virus titration of the main samples.

The testing was performed with the *Transmissible Gastroenteritis Virus of Swine (TGEV)* which served as a model virus for the *Middle East Respiratory Syndrome Coronavirus (MERS-CoV)*.

**Information about the testing**

<i>Principal:</i>	PMF Natural Products company	<i>Test run:</i>	S3
<i>Product(s):</i>	PMF-concentrate	<i>Test date:</i>	12.03.2015
<i>Test system:</i>	TGEV ( <i>Toyama</i> ) + ST75/2-cells	<i>Analysis:</i>	18.03.2015 (6 p.i.)

**Test methodology and test parameters**

*Test method:* quantitative virucidal suspensions test according to DVV/RKI-guideline (*Version 08/08*)  
*Test mixture:* 1 VT protein load + 1 VT virus suspension + 8 VT 1,25fold working solution  
*Protein load:* no additional protein load (*PBS*)  
*Parameter:* test temperature: 37° C with the exposure time(s) of: 60 and 240 min.

**Tested product sample(s)**

*1<sup>st</sup> product:* PMF-concentrate [*Product sample: as received (designation: PMF), Arrival: 01.08.2014, Storage at 2-8° C*]

**Tab. 1: Weight of content**

Set	Product(s)	Conc. in Test (1x)	Working sol. (x 1,25)	Dosage	pH of working sol.
#1	PMF-concentrate	20,34%	25,42%	0,75 g in 2,95 mL	pH 9,71 (in test: pH 9,69)

**Tab. 2: Content of samples**

Samples	1a	1b	2a	2b
	Virus inactivation			
	Set #1 / 60 min.		Set #1 / 240 min.	
PBS	15µL	15µL	15µL	15µL
TGEV	15µL	15µL	15µL	15µL
PMF / Sol.	120µL	120µL	120µL	120µL
Titration	<i>Lycke (VF = 1000)</i>		<i>Lycke (VF = 1000)</i>	

Samples	3a	3b	4a	4b	5
	Virus control / 60 Min.		Virus control / 240 Min.		Cytotoxicity
	w/o		w/o		Set #1/240 min.
PBS	15µL	15µL	15µL	15µL	15µL
TGEV	15µL	15µL	15µL	15µL	
Medium					15µL
PBS	120µL	120µL	120µL	120µL	
PMF / Sol.					120µL
Titration	S&K (VF = 5)		S&K (VF = 5)		<i>Lycke (VF = 1000)</i>

**Performing of the test**

**1. Preparation of the product solution:** (in the specified sequence)

- **0,75 g PMF-concentrate** was solved with agitation and warming to 37°C in 2,95 mL A. bidest.

**2. Preparation of the test samples**

- Per test point (concentration/exposure time) 2 redundant test samples were prepared.
- Test mixture: 1 vol. PBS + 1 vol. virus suspension + 8 vol. PMF-working solution (1,25-fold)

**3. Dilution of the test sample and estimation of virus titer**

- **Termination of virus inactivation:** after exposure the test samples were diluted with medium (cf. virus titration).
- With the **virus control** the virus titer was estimated using the methodology of *Spearman & Kärber* with VF = 5 from 113 µL (out of 150 µL of the test sample).
- With the **virus inactivation samples** the virus titer was estimated using the methodology of *Lycke*. For each of the test samples (a and b) 48 µL was added to 96 mL Medium, corresponding to a dilution of VF = 1000. All of the 96 mL were then transferred to cell cultures with 200 µL per well (480 wells).

**4. Susceptibility control**

- Sample 5 (cytotoxicity sample) was diluted 1000fold and was then distributed to cell cultures (cf. virus inactivation). Afterwards a virus dilution serie (VK/E) was transferred to these cells.

**5. Judgement of the cells / virus detection**

- At day 6 p.i. the cell cultures were examined visually using a microscope (magnification: 100fold). The virus positive cell cultures were identified by the virus induced CPE).

**Tab. 3.1: Virus control + Susceptibility control** (virus titration: according to *Spearman & Kärber*)

Samples	3a	3b	Ø	4a	4b	Ø	5	VK/E
	Virus control / 60 min.			Virus control / 240 min.			Susceptibility Control	
1 / -0,7	4/4 <sup>1</sup>	4/4	<b>8/8</b>	4/4 <sup>1</sup>	4/4	<b>8/8</b>	8/8	8/8
2 / -1,4	4/4	4/4	<b>8/8</b>	4/4	4/4	<b>8/8</b>	8/8	8/8
3 / -2,1	4/4	4/4	<b>8/8</b>	4/4	4/4	<b>8/8</b>	8/8	8/8
4 / -2,8	4/4	4/4	<b>8/8</b>	4/4	4/4	<b>8/8</b>	8/8	8/8
5 / -3,5	4/4	4/4	<b>8/8</b>	4/4	4/4	<b>8/8</b>	8/8	8/8
6 / -4,2	4/4	4/4	<b>8/8</b>	4/4	4/4	<b>8/8</b>	8/8	8/8
7 / -4,9	4/4	4/4	<b>8/8</b>	4/4	3/4	<b>7/8</b>	8/8	8/8
8 / -5,6	3/4	4/4	<b>7/8</b>	2/4	2/4	<b>4/8</b>	4/8	3/8
9 / -6,3	1/4	1/4	<b>2/8</b>	0/4	0/4	<b>0/8</b>	0/8	2/8
10 / -7,0	1/4	0/4	<b>1/8</b>					1/8
11 / -7,7	0/4		<b>0/8</b>					0/8
ZK	0/4	0/4	<b>0/8</b>	0/4	0/4	<b>0/8</b>	0/8	0/8
Titer/test vol. (lg ID <sub>50</sub> )	6,13	6,13	<b>6,13</b>	5,6	5,43	<b>5,52</b>	5,78 ± 0,39	5,60 ± 0,26
Average ± CI (95%) <sup>2</sup>	6,13 ± 0,34 per 100 µL (≈ 6,11 lg ID <sub>50</sub> pro 96 µL)			5,52 ± 0,32 per 100 µL (≈ 5,50 lg ID <sub>50</sub> pro 96 µL)			RF = 0,18 ± 0,47	
<b>Reduction</b> <sup>3</sup> lg ID <sub>50</sub> ± CI [95%]	-			<b>0,61 ± 0,46</b>			<b>cells susceptible: yes</b> <sup>4</sup>	

<sup>1</sup> = number of virus positive cell culture units to total number of cell culture units

<sup>2</sup> = Calculation of 95% confidential intervall of virus titer as well as virus reduction following DVV/RKI-Guideline.

<sup>3</sup> = Virus reduction: titer of virus control minus titer of sample (lg ID<sub>50</sub>).

<sup>4</sup> = Susceptibility of the detection cells is to be assumed when Δ virus titer is ≤ lg 0,5 [DVV/RKI-Guideline].

**Tab. 3.2: Virus inactivation** (virus titration: according to Lycke)

Samples	1a + 1b	2a + 2b
	Virus inactivation (VF = 1000)	
	Set #1 / 60 min.	Set #1 / 240 min.
analysed sample vol.	2 x 48 = 96 µL	2 x 48 = 96 µL
Cell culture units	480	480
Virus positive	43	0
Ratio p <sup>2</sup>	0,0896	0,0
Residual virus (lg ID <sub>50</sub> per 96 µL)	1,80 ± 0,34	< 1 ID <sub>50</sub> (0,0 lg ID <sub>50</sub> )
Virus input (lg ID <sub>50</sub> per 96 µL)	6,11 ± 0,34	5,50 ± 0,32
<b>Reduction<sup>3</sup></b> (lg ID <sub>50</sub> ± CI [95%])	<b>4,31 ± 0,48</b>	<b>&gt; 5,50 ± 0,45</b>

<sup>1</sup> = sample volume transferred onto cell cultures: 48 µL from test mix a. plus 48 µL from test mix b. resulting in 2 x 48 = 96 µL

<sup>2</sup> = ratio of virus positive cell culture units to total number of cell cultures.

<sup>3</sup> = Virus reduction: titer of virus control (cf. Tab. 3.1) minus titer of sample (lg ID<sub>50</sub>)

***Estimation of virus titer by LYCKE's method*** (Arch Ges Virusforsch (1957); 7:483-493)

Calculation of virus titer by using the following formula:

$$- \text{ID}_{50} = [1,4 \times \ln (1-p)] \quad p = \text{ratio of positive cell cultures to total number of cell cultures}$$

- **Example:** 51 out of 100 cell culture units was virus positive →  $p = 51/100 = 0,51$

p put into formula:                    -  $\text{ID}_{50} = [1,4 \times \ln (1- 0,51)]$

with  $\ln (0,49)$ :                         -  $\text{ID}_{50} = 1,4 \times -0,71$

resulting in:                              -  $\text{ID}_{50} = -0,998$  or  $\text{ID}_{50} = 0,998$

That means that per single cell culture unit 0,998 or 1 ID<sub>50</sub> of residual virus was present.

When this content of virus was multiplied with the number of cell culture units (= 100) the complete amount of residual virus was obtained:  $1,0 \text{ ID}_{50} \times 100 = 100 \text{ ID}_{50}$  or  $\lg \text{ID}_{50} = 2,0$

**Result of the example:** the total quantity of residual virus which was present in the examined sample of liquid was estimated to **lg ID<sub>50</sub> = 2,0**



**Materials and reagents used:**

• **Testvirus**

Test virus	Transmissible Gastroenteritis Virus of Swine (TGEV)
Strain	Toyama 36
Origin	Virusbank der BFA f. Viruskrankheiten der Tiere; Friedrich Löffler-Institut, Insel Riems Virus (lyophilisate) v. 05/2003; kindly provided by Dr. M. Dauber (Virus passage FLI +0)
Virus material used in test	Supernatant from infected cell culture, Virus propagation TGEV-12 v. 16.02.2015; Set #1 Virus passage: FLI +12;

• **Cells**

Cells	ST75/2 cells (foetal testis cells of swine)
Origin	Robert Koch-Institut, Berlin Cells received 03/2002 in frozen condition (1 Ampoule; v. 13.02.1996); corresponding to cell passage RKI +0
Cell passage used in test	RKI + 3 / + 15

• **Additional material and reagents**

<i>Material</i>	<i>Supplier</i>	<i>Order No.</i>	<i>Lot</i>	<i>Expiry date</i>
DMEM	Biochrom	F 0435	1006 C	11/2015
Glutamine	Biochrom	K 0283	0978 B	08/2016
Pen./Strept.	Biochrom	A 2213	0627 C	05/2017
FCS	Biochrom	S 0210	0677 B	06/2019
PBS	Biochrom	L 1820	0743 C	07/2017
Trypsin	Invitrogen	25300-096	1437736	09/2015

• **Performing of the experiment and responsibilities**

Part of experiment	Performed by (position)
Supervision	Dr. Ch. Jursch (Laborleiter)
Control of product input	Fr. S. Sachs (Biologielaborantin) und Dr. Ch. Jursch (Laborleiter)
Performing the test	Fr. S. Sachs (Biologielaborantin)
Cell culturing	Fr. S. Sachs (Biologielaborantin)
Reading of cells & Raw data	Dr. Ch. Jursch (Laborleiter)
Data input & Analysis	Dr. Ch. Jursch (Laborleiter)
Protocol preparation	Dr. Ch. Jursch (Laborleiter)