

## ACCUMULATION OF TRACE METALS IN MANGROVE PLANT *SONERATIA CASEORALIS* IN SONGKHLA LAKE, THAILAND

PRADIT, S.<sup>1\*</sup> – SHAZILI, N. A. M.<sup>2</sup> – PATTARATUMRONG, M. S.<sup>1</sup> – CHOTIKARN, P.<sup>1</sup> –  
KOBKEATTHAWIN, T.<sup>1</sup> – YUCHAROEN, M.<sup>1</sup> – TOWATANA, P.<sup>1</sup>

<sup>1</sup>*Coastal Oceanography and Climate Change Research Center, Marine and Coastal Resources  
Institution, Prince of Songkla University, Songkhla Province, Thailand*

<sup>2</sup>*Ocean Research Institute, Univerisiti Malaysia Terengganu, Kuala Terengganu, Malaysia*

*\*Corresponding author*

*e-mail: spradit22@gmail.com; phone: +66-74-282-329; fax: +66-74-212-782*

(Received 19<sup>th</sup> Apr 2018; accepted 13<sup>th</sup> Jun 2018)

**Abstract.** Mangrove sediment and mangrove plant *Sonneratia caseolaris* in Songkhla Lake, Thailand have been increasingly threatened by trace metal pollution. Mangroves receive trace metal pollution from upstream areas and the sea. However, little is known about the mangrove plant's capacity to uptake and store trace metals. In this study, the concentrations of As, Cd, Pb and Zn in mangrove plant parts (leaf, root, and bark) of *Sonneratia caseolaris* and sediments were determined. Sediment and plant samples were digested using total metal concentration procedure and ICPMS techniques. It was found that the maximum values of total concentrations of As in mangrove sediments were classified as heavily polluted according to USEPA. The highest concentrations of As and Zn were found to be in bark whereas Cd and Pb in the root. The fractions of all the metals were measured using BCR techniques and their results revealed that the bioavailability fraction was greater than the residual fraction. The order of the amounts of trace metals present in the fractions was as follows; As: Oxidizable > Reducible > Residual > Exchangeable and the order of Cd: Exchangeable > Reducible > Oxidizable > Residual. Both BCF and TF values showed limited accumulation of these elements in their aboveground parts and thus presented a low food chain hazard except at Phawong and U-Taphao canals.

**Keywords:** *BCR techniques, coastal lagoon, BCF, TF, food chain*

### Introduction

Mangrove forest is a group of trees and shrubs and commonly found in coastal areas. Mangrove is like a connection place between land and sea and important to the ecosystem especially to the aquatic population. Mangroves are an important food source of aquatic fauna in the tidal areas. In addition, the elongated root system makes it possible to trap sediment and trap pollutants such as trace metals. Generally, mangrove increases metal accumulation in sediments by modifying the soil acidity, redox potential, organic contents and salinity (Zhou et al., 2010; Sekomo et al., 2011) and subsequently reduces metal exposure to adjacent aquatic environment (Nath et al., 2013). Active uptake requires metabolic energy and takes place against a chemical gradient. The rate of heavy metals uptake is corresponding with its available pool at the root surface (Lanno et al., 2004). For example, when As, Cd and Pb are present in soluble forms in pore water, plant roots are able to take up great amounts of these metals. The uptake rates increase as concentration in the solutions increase. The plant uptake is also affected by temperature. According to Pendias and Pendias (2001), a higher ambient temperature influences a greater uptake of trace metals by plants.

It is widely recognized that the adverse impact of heavy metals does not simply depend on their concentrations but critically on their bioavailable fraction of the total metal

concentration in sediment (Kim et al., 2015; Tokalioglu et al., 2000). Toxicological bioavailability is defined as a bioaccumulation effect of heavy metals within plants. Among various pollutants, trace metals with persistence, non-biodegradation, toxicity and bioavailability pose a major threat to the mangrove biodiversity and human health (Liu et al., 2014). Mangroves have the capacity to contain trace metals by accumulating them in their sediments and plant tissues (MacFarlane et al., 2003). Accumulation of these metals also can cause potential problems to animals and human through food chain interaction (Hosani and Anouti, 2014).

*Sonneratia caseolaris* is one of the mangrove trees found in Songkhla lake, Thailand and a lead or pioneer species. In the Outer Section of Songkhla Lake, *Sonneratia caseolaris* is generally found at the mouth of the Phawong canal and U-Taphao canal and Kuannieng District, the Middle Section of Songkhla Lake is found in Kukhut non hunting areas as well. In the Songkhla Lake, there were several studies on trace metal contamination in the mangrove sediment but little is known about heavy metal uptake by mangrove plant (*Sonneratia caseolaris*). Therefore, the aims of this study were to determine the bioavailability of As, Cd, Pb and Zn in mangrove sediments of the Songkhla Lake using modified European Community Bureau of Reference (BCR) sequential extraction procedure and to analyze the accumulation of trace metals (As, Cd, Pb and Zn) in mangrove sediment and mangrove plant parts (leaves, root, bark) of *Sonneratia caseolaris*.

## Material and method

### Study area

The Songkhla Lake is a shallow coastal lagoon located in the southern part of Thailand and occupies approximately 1,042 km<sup>2</sup> formed by an interaction of land and ocean processes over geological time (Sompongchaiyakul and Sirinawin, 2007). The lake is divided into 4 sections; Thale Noi, Inner Section, Middle Section and Outer Section. The salinity ranges from fresh water in Thale Noi to saline water in the Outer Section. The Outer Section is connected to the Gulf of Thailand through a deep and narrow outlet at Songkhla city. The system receives runoff and wastewater from the surrounding watershed. The runoff carries sediment onto the lakes, which will be transported through the lake by the general movement of water towards the Gulf of Thailand. According to the EmSong Project (1998), the average residence time of the water mass in the Inner, Middle and Outer Lakes is 55, 28 and 15 days respectively. The average sediment accumulation rate of in the lake is 2.5-3.5 mm/year and at some places it is higher than 15 mm/year (Sojisuporn, 2005). The average depth of Inner, Middle and Outer Lakes is 2, 1.5 and 1.5 meters, respectively. The lake is facing problems of environmental degradation due to urban expansion and industrialization. Sources of pollutants to the lake system include municipal wastes from Hat Yai and Songkhla cities, industrial wastes mainly related to the rubber industries, seafood processing industries, mining activities and pollution from boats in Songkhla harbour (Sompongchaiyakul and Sirinawin, 2007; Pradit et al., 2010).


### Sample collection and preparation

The sediment cores were collected from 4 stations (*Fig. 1*) which were located in the Middle Section (station 1: Kukhut non hunting area) and in the Outer Section (station 2: Kuannieng, station 3: Phawong canal and station 4: U-Taphao canal). The selected station

based on the appearance of *Sonneratia caseolaris* was found in the Middle Section (Kukhut non hunting area; station 1) and in the Outer Section of the lake (Kuannieng, Phawong and U-Taphao canals). The survey was carried out in November, 2016.

The surface of mangrove sediment (0-10 cm) was taken from the active root zone using a Plexiglas tube (5 cm diameter and 30 cm length) by pushing into the bottom sediment with 3 replicates for each station. After that the sediment cores were sectioned 5 cm (1 core got 2 pieces) and then packed into a plastic zip lock, labeled and put in to a plastic container before being transported to the laboratory.

Mangrove plant; *Sonneratia caseolaris* samples consisted of root, bark and leaf. The leaves (old leaf = yellow colour, mature leaf = green colour and young leaves) were collected by hand pick (in total about 30 leaves per stations) whereas roots (about 10 roots per station) and barks (about 5 pieces; each piece about 3 × 5 cm) were collected using a ceramic knife. The plant samples were stored in clean plastic bags. The 200 g wet sample (plant and sediment) of each station was put on tray and then oven dried at 50 °C, homogenized and grinded in a mortar and sieved through a 1 mm nylon sieve and then stored in plastic zip locked bags with label and kept in desiccators prior to further analysis.



**Figure 1.** Map of sampling station in Songkhla Lake (station 1: Kukhut non hunting area; station 2: Kuannieng; station 3: Phawong canal; station 4: U-Taphao canal)

### Sample analysis

The bulk sediment samples (about 0.1 g) were totally digested in an acid mixture (5 ml HCl: 8 ml HNO<sub>3</sub>: 2 ml HF) and analyzed for trace metals, according to the published methodologies of Noriki et al. (1980) Chojnacka et al. (2004) and Loring and Rantala (1992), with some modifications (the Teflon vessels were heated in an oven instead of a

microwave until complete digestion of the sample occurred). The amounts of trace metals (As, Cd, Pb and Zn) were measured by ICP-MS (Perkin Elmer Elan 9000). The sediment certified reference material; SRM 1646/estuarine sediment was similarly analyzed to validate the accuracy of the analytical procedure. The analytical values were within 90 % of the certified values, which demonstrated the validity of the methods applied. Plant sample was digested in a closed teflon vessel. Sample of 0.5 g was dissolved in 5 ml of HNO<sub>3</sub> (Chojnacka et al., 2004). Concentrations of As, Cd, Pb and Zn in digested samples were then determined by Inductively Coupled Plasma Mass Spectrometry (ICPMS). Standard reference material for plant (Peach Leaves: NIST 1547) was digested following the same procedure and analyzed for metal concentrations to validate accuracy and precision of the analysis.

Sequential extraction of mangrove sediment was performed using the revised BCR protocol (Rauret et al., 1999) briefly describes as follows:

Step 1 (Fraction 1: Exchangeable and acid soluble fractions): 40 ml of 0.11 mol/l acetic acid was added to 1 g of the air-dried sediment sample in a 50 ml polyethylene centrifuge tube. The tube was shaken for 16 h at room temperature at a speed of 23 ± 1 rpm. The extract was separated from the solid residue by centrifugation (3000 rpm for 20 min), decanted into a polyethylene bottle and stored at 4 °C. The residue was washed by shaking with 20 ml of distilled water for 15 min and centrifuging after which the supernatant was discarded.

Step 2 (Fraction 2: Reducible fraction): 40 ml of 0.5 mol/l hydroxylamine hydrochloride (adjusted to pH 1.5 by addition of a fixed amount of HNO<sub>3</sub>) was added to the residue from step 1, and the extraction performed as in step 1 above.

Step 3 (Fraction 3: Oxidizable fraction): 10 ml of 8.8 mol/l hydrogen peroxide was added in aliquots to the residue from step 2. The vessel was covered loosely and the contents were digested at room temperature for 1 h with occasional agitation. It was then placed in a water bath and digested at 85 °C until the volume was reduced to less than 3 ml. Another 10 ml of the hydrogen peroxide was added, and further heated to near dryness. Thereafter, 50 ml of 1.0 mol/l ammonium acetate (adjusted to pH 2 with HNO<sub>3</sub>) was added, and the extraction was performed as in the previous steps.

Step 4 (Fraction 4: Residual fraction): The residue from step 3 was transferred into a suitable vessel and the metal content was determined by microwave-assisted digestion with *aqua regia*. This step was for analyzing the metals content in their primary and secondary minerals.

The trace element (As, Cd and Pb) content of each fraction was analyzed using ICP-MS model Perkin Elmer Elan 9000.

### ***Data analysis and statistical***

Two comparative measures were chosen to assess the trace metals uptake and distribution within the plant. Bioconcentration factor (BCF) and translocation factor (TF) was calculated using the following equations:

$$\text{BCF} = [\text{plant tissue}] / [\text{sediment}]$$

$$\text{TF} = [\text{shoot}] / [\text{root}] \quad (\text{TF} > 1: \text{high mobility})$$

where [ ] is metal concentration in µg/g.

In order to evaluate the degree of trace metal amount in the mangrove sediments of mangrove ecosystem, ecological risk assessment was conducted using the Håkanson, ecological risk index ( $R_I$ ) (Eqs. 1 and 2; Håkanson, 1980).

$$R_I = \sum Er^i = \sum Tr^i Cf^i \quad (\text{Eq.1})$$

$$Cf^i = C_o^i / C_n^i \quad (\text{Eq.2})$$

where:

$R_I$  = the sum of all risk factors for heavy metals in sediments

$Er^i$  = the monomial potential ecological risk factor

$Tr^i$  = the toxicity coefficient which represents the toxic response factor for a given metal. The value of Tr for As, Cd, Pb and Zn is 10, 30, 5 and 1 respectively

$Cf^i$  = the contamination factor

$C_o^i$  = the concentration of metal in the sediment of mangrove ecosystem

$C_n^i$  = the background value of heavy metal in coastal sediment

### Statistic analysis

The data analysis using ANOVA to detect if any significant difference in mean existed between the mangrove sediment and plant parts. Duncan's new multiple range test (DMRT) was used to describe the differences between the variables (sediment and plant parts). Significance was set at 95% confidence level. Correlation was used to clarify the relationship between sediment fractions and metal accumulation in plant parts.

## Results

### Sediment characteristic in mangrove sediment

The metal concentrations in sediments from of Songkhla Lake (Table 1) ranged from 20.4 (station 1) - 50.5 (station 2)  $\mu\text{g/g}$  dry weight for As, 0.006 (station 4) - 0.182 (station 3)  $\mu\text{g/g}$  dry weight, for Cd, 48.8 (station 4) - 78.8 (station 1)  $\mu\text{g/g}$  dry weight, for Pb and 48.6 (station 4) - 126.6 (station 1)  $\mu\text{g/g}$  dry weight, for Zn. The amounts of As, Zn and Pb were highest at Kuannieng (station 2) whereas Cd was highest at Phawong canal (station 3). The amounts of Zn, Cd and Pb were lowest at mouth of U-Taphao canal (station 4) whereas As was lowest at Kukhut (station 1). Besides that the results of the selected physical properties of sediment and water samples were summarized in Table 1. The pH in water ranged from 7.03 – 7.85 and in sediment ranged from 6.71-7.53. Salinity ranged between 8 ppt (station1) – 20 ppt (station 2).

**Table 1.** Trace metals, physical properties of sediment and water in Songkhla Lake

Station	pH water	Salinity (ppt)	pH sed	Eh (mV)	As	Zn	Cd	Pb
					$\mu\text{g/g}$ dry weight (in sed.)			
St.1	7.33	8	7.53	-292	20.35	86.5	0.0759	59.8
St.2	7.85	20	6.71	-63	50.5	126.6	0.0984	78.8
St. 3	7.42	17	7.4	-149	28.7	102.1	0.1818	51.3
St.4	7.03	10	6.99	-79	31.7	48.6	0.0061	48.8

### Accumulation of trace metals in *Sonneratia caseolaris*

Accumulation of trace metals (As, Cd, Pb and Zn) in *Sonneratia caseolaris* is shown in Table 2 and summarized as below:

As: The amount of As ranged from 0.20-4.72 µg/g dry weight. It was found that the concentration of As was highest in the barks (4.72 µg/g dry weight) at U-Taphao Canal (station 4), followed by roots (1.51 µg/g dry weight) and old leaves (1.03 µg/g dry weight).

Cd: The amount of Cd ranged from 0.003-0.041 µg/g dry weight. The result revealed that the concentration of Cd was highest in root (0.041 µg/g dry wt.) at Phawong canal (station 3), followed by old leaves (0.033 µg/g dry weight), green leaves (0.029 µg/g dry weight) and barks (0.029 µg/g dry weight).

Pb: The amount of Pb ranged from 0.18-6.76 µg/g dry weight. It was found that the concentration of Pb was highest in the root (6.76 µg/g dry wt.) at U-Taphao canal (station 4), followed by bark (4.12 µg/g dry weight) and old leaves (2.74 µg/g dry weight).

Zn: The amount of Zn ranged from 2.66-35.80 µg/g dry weight. The result showed that concentration of Zn was highest in the bark (35.80 µg/g dry wt.) at Kuannieng (station 2), followed by young leaves (17.10 µg/g dry weight) and root (16.58 µg/g dry weight).

Overall the amounts of trace metals were in order Zn > Pb > As > Cd.


**Table 2.** Trace metals content in different parts of *Sonneratia caseolaris* of Songkhla Lake

Station	Plant part	Zn	As	Cd	Pb
		(µg/g dry weight)			
1	Root	16.16	0.83	0.021	1.39
1	Bark	9.90	0.20	0.029	0.94
1	Leaf_Y	17.10	0.26	0.009	0.28
1	Leaf_M	12.98	0.29	0.014	0.48
1	Leaf_O	12.08	0.70	0.022	0.73
2	Root	24.40	1.51	0.027	1.85
2	Bark	35.80	0.79	0.026	2.08
2	Leaf_Y	4.14	0.04	0.006	0.18
2	Leaf_M	11.10	0.36	0.029	1.12
2	Leaf_O	10.44	0.57	0.033	2.74
3	Root	16.58	0.89	0.041	2.12
3	Bark	2.66	0.28	0.010	0.43
3	Leaf_Y	11.20	0.26	0.004	0.27
3	Leaf_M	5.36	0.53	0.006	0.44
3	Leaf_O	2.84	0.95	0.007	0.32
4	Root	5.88	0.41	0.003	6.76
4	Bark	8.04	4.72	0.019	4.12
4	Leaf_Y	7.60	0.37	0.004	0.21
4	Leaf_M	4.80	0.72	0.005	0.26
4	Leaf_O	4.76	1.03	0.006	0.70

Remarks: Y = young leave, M = mature leave (green colour), O = old leave (yellow colour)

### Trace metal speciation in mangrove sediment

The speciation of As, Cd and Pb in the mangrove sediment samples from the Songkhla Lake using the BCR protocol were presented in *Figure 2*.



**Figure 2.** The speciation distribution of As, Cd and Pb in mangrove sediments in Songkhla Lake, Southern Thailand

The results of the sequential extraction for overall stations were as follows:

Exchangeable and acid soluble fraction (F1): Cd (49.58% - 64.33%) was found dominant in the acid-soluble fraction (F1), associated with exchangeable cations and carbonates for all the stations.

Reducible Fraction (F2): Pb (40.89% - 74.81%) was found dominant in F2 in all the stations except station 3. This fraction is readily available with medium mobility and may be released into the environment under unstable anoxic conditions.

Oxidizable fraction (F3): As (32.47% - 64.76%) was found dominant in F3 at all the stations except station 4. Metals bound to organic matter are reasonable stable in nature however under strong oxidizing conditions such as currents, dredging, flooding and tides the organic matter can be degraded, hence leading to a release of heavy metals bound to this component (Sarkar et al., 2014; MacFarlane and Burchett, 2000).

Residual fraction (F4): Pb (2.60% - 16.49%) was found highest in F4. The residual solid normally consists of mainly primary and secondary minerals that retained heavy metals within their crystal structure. They are regarded as immobile and unavailable fraction.

As: Oxidizable (F3) > Reducible (F2) > Residual (F4) > Exchangeable (F1)

Cd: Exchangeable (F1) > Reducible (F2) > Oxidizable (F3) > Residual (F4)

Pb: Reducible (F2) > Oxidizable (F3) > Residual (F4) > Exchangeable (F1)

### **Bioaccumulation factors (BAF) and transfer factors (TF)**

The bioconcentration factor (BCF) of *Sonneratia caseolaris* in different plant parts was varied as As>Cd>Zn> Pb. The BAF of Zn, As, Cd and Pb was shown in Table 3. The bioconcentration factor (BCF) was always below 1 for all the trace metals and all the stations except As (1.06) in old leaf at Phawong canal (station 3) and Cd in old leaf (1.02) at the U-Taphao canal (station 4). These results suggested that *Sonneratia caseolaris* showed the limited uptake of Zn and Pb (BCF < 1) but surprisingly not for As and Cd which BCF > 1 in old leaf was found in Phawong canal (station 3) and U-Taphao canal (station 4).

**Table 3.** Bioconcentration factor (BCF) of Zn, As, Cd and Pb in *Sonneratia caseolaris* of Songkhla Lake

Station	Plant part	Zn	As	Cd	Pb
1	Root	0.19	0.04	0.27	0.02
1	Bark	0.11	0.01	0.38	0.02
1	Leaf_Y	0.20	0.01	0.11	0.00
1	Leaf_M	0.15	0.01	0.19	0.01
1	Leaf_O	0.14	0.03	0.28	0.01
2	Root	0.19	0.03	0.27	0.02
2	Bark	0.28	0.02	0.26	0.03
2	Leaf_Y	0.03	0.00	0.06	0.00
2	Leaf_M	0.09	0.01	0.29	0.01
2	Leaf_O	0.08	0.01	0.34	0.03
3	Root	0.16	0.03	0.23	0.04
3	Bark	0.03	0.01	0.06	0.01
3	Leaf_Y	0.11	0.01	0.02	0.01
3	Leaf_M	0.05	0.02	0.03	0.01
3	Leaf_O	0.17	1.06	0.16	0.15
4	Root	0.12	0.01	0.42	0.14
4	Bark	0.17	0.01	3.19	0.08
4	Leaf_Y	0.16	0.01	0.70	0.00
4	Leaf_M	0.10	0.02	0.74	0.01
4	Leaf_O	0.10	0.03	1.02	0.01

Y = young leave, M = mature leave (green colour), O = old leave (yellow colour)



The translocation factor (TF) of *Sonneratia caseolaris* in different plant parts varied as Cd >As>Zn> Pb (Table 4). The TF value of Zn ranged 0.16 -1.47 µg/g dry weight; As ranged 0.02-2.55 µg/g dry weight; Cd ranged 0.10-7.60 µg/g dry weight; Pb ranged 0.03-1.48 µg/g dry weight. The TF of these metals was above 1 at some stations and below 1 at other stations. At U-Taphao canal (station 4) TF value was above 1 for Zn, As, and Cd. TF value of As and Cd in old leaf was greater than one at station 4 and 1. TF value of Pb in old leaf was greater than one at station 2. TF value in bark was greater than 1 for Zn at station 1 and 4; As at station 4; Cd at station 1 and 4; Pb at station 2. Thus the result revealed that this plant species was effectively transferring Cd to the aerial parts. The TF of Pb was always below 1 (except station 2 in bark and station 3 in old leaf), which might be described as having moderate mobility for this metal.

**Table 4.** Translocation factor (TF) for mangrove plant parts

Station	Plant part	Zn	As	Cd	Pb
1	Bark	0.61	0.24	1.39	0.68
1	Leaf_Y	1.06	0.31	0.42	0.20
1	Leaf_M	0.80	0.35	0.70	0.35
1	Leaf_O	0.75	0.85	1.05	0.53
2	Bark	1.47	0.52	0.96	1.12
2	Leaf_Y	0.17	0.02	0.22	0.10
2	Leaf_M	0.45	0.24	1.08	0.60
2	Leaf_O	0.43	0.37	1.23	1.48
3	Bark	0.16	0.31	0.24	0.20
3	Leaf_Y	0.68	0.29	0.10	0.13
3	Leaf_M	0.32	0.60	0.14	0.21
3	Leaf_O	0.17	1.06	0.16	0.15
4	Bark	1.37	1.16	7.60	0.61
4	Leaf_Y	1.29	0.90	1.67	0.03
4	Leaf_M	0.82	1.78	1.76	0.04
4	Leaf_O	0.81	2.55	2.42	0.10

Y = young leave, M = mature leave (green colour), O = old leave (yellow colour)

### **Ecological risk index of heavy metals in mangrove sediments of different areas**

The potential ecological risk index of single element  $E_r^i$  shown in Table 5 indicated that As and Zn were classified as moderate level whereas Cd and Pb were classified as low level. As shown in Table 5, the ecological risk indexes ( $R_1$ ) of trace metals in the mangrove sediments from different part of Songkhla Lake showed that Kuannieng (station 2) and Phawong canal (station 3) were classified as moderate level whereas Kukhut nonhunting area (station 1) and U-Taphao canal (station 4) were classified as low level.

### **Comparison of trace metal concentration with Sediment Quality Guidelines (SQGs)**

Several sediment quality guidelines (SQGs) for the assessment of sediment quality using chemical and biological effect databases have been established. These SQGs are

summarized in Table 6. NOAA presents ERL (effects range low) and ERM (effects range mean) guidelines for estuarine and marine environments which represent the 10<sup>th</sup> and 50<sup>th</sup> percentiles of adverse biological effects (NOAA, 1999). The MacDonald et al. (2000) SQGs for freshwater environments have a lower TEC (threshold effects concentration) and an upper PEC (probable effect concentration) at which toxicity to bottom dwelling organisms are predicted to be unlikely and probable, respectively. The US-EPA has also made classifications (non-polluted, moderately polluted, heavily polluted based on toxicity tests (Baudo et al., 1990; Filgueiras et al., 2004). Regional background values assessed from core sediment samples were reported by Choi et al. (2008) and offshore sediments in the Gulf of Thailand by Shazili et al. (1999). Background levels varied from 55-115 µg/ g dry weight for Zn, from 5-9 µg/g dry weight for As, from 0.03-0.2 µg/g dry weight for Cd, from 15-30 µg/g dry weight for Pb.

**Table 5.** Ecological risk index of trace metals in mangrove sediments of Songkhla Lake

$E_r^i$	Potential ecological risk for single regulator		$R_I$	Ecological risk for all factors	
<40	Low		$R_I < 95$	Low (L)	
$40 \leq E_r^i \leq 80$	<b>Moderate</b>		$95 \leq R_I \leq 190$	Moderate (M)	
$80 \leq E_r^i \leq 160$	Considerable		$190 \leq R_I < 380$	Considerable (C)	
$160 \leq E_r^i \leq 320$	High		$R_I \leq 380$	Very high (V)	
$E_r^i \geq 320$	Very high				
Sites	$E_r^i$			$R_I$	
	Zn	As	Cd	Pb	
1 Kukhut	<b>44.0</b>	29.1	3.2	2.7	78.9 (L)
2 Kuannieng	<b>64.4</b>	<b>72.1</b>	4.1	3.5	144.1(M)
3 Phawong canal	<b>51.9</b>	<b>41.0</b>	7.6	2.3	102.8 (M)
4 U-Taphaocanal	24.7	<b>45.3</b>	0.3	2.2	72.4 (L)

**Table 6.** Comparison the metal concentration (µg/g dry weight) and the Sediment Quality Guidelines (SQGs)

Element	US EPA			NOAA		MacDonald et. al. (2000)		This study
	Non polluted	Moderately polluted	Heavily polluted	ERL	ERM	TEC	PEC	
Pb	<40	40-60	>60	46.7	218	36	130	48.8-78.8
As	<3	3-8	>8	8.2	70	9.8	33	20.35-50.5
Cd			>6	1.2	9.6	0.99	5	0.0061-0.1818
Zn	<90	90-200	>200	150	410	120	460	48.6-126.6

TEC: threshold effect concentration, PEC: probable effect concentration, ERL: effects range low, ERM: effects range mean

Sediment quality guidelines (SQGs) were recognized as appropriate thresholds to reveal adverse effects of heavy metals in sediments for plants, animals and human health. SQGs were applied to our study to determine the overall pollution status in sediments of Songkhla Lake. The amount of As and Pb was above the background value. Our study showed that highest amount of As at Kuannieng (station 2) of

Songkhla Lake exceeded the ERL value; TEC value was classified as heavily polluted probably affected the health of plants and organisms. Pb levels from both sites exceeded the ERL and TEC value and were regarded as heavily polluted at Kuannieng (station 2) and classified as moderately polluted for stations 1, 3, and 4. Cd level was classified as lower than ERL and TEC. Zn was classified as moderately polluted and exceeded TEC value.

***The comparison of trace metals in mangrove sediment and plant parts***

Our result revealed that the total concentration of all metals (As, Zn, Pb, Cd) in mangrove sediment no statistically significant difference among the stations ( $p > 0.05$ ). For the comparison of the trace metal concentrations between plant parts of *Soneratia caseolaris*, all the values show variations among the stations. The post hoc Duncan-Test showed no significantly different accumulation in barks ( $p > 0.05$ ) whereas roots and leaves were significantly different ( $p < 0.05$ ). The comparison of the metal contamination in the different sediment fractions revealed the low metal concentration in residual fraction and high in bioavailability fractions. The post hoc Duncan-Test showed among the stations showed no significant difference in residual fraction ( $p > 0.05$ ) whereas the other fractions (Exchangeable, Reducible and Oxidizable) were significantly different ( $p < 0.05$ ).

***Correlation between fractionation of heavy metals in sediment and accumulation in plant parts***

The accumulation in various parts of *Soneratia caseolaris* was investigated to determine the relationship between the fractionation and plant's uptake as showed in Table 7. Fraction 1 (Exchangeable) was significantly negative correlated with accumulation in leaves (young, green, old). The result indicated that Fraction 2 (Reducible) was showed positive correlation with root ( $r = 0.634$ ,  $p < 0.05$ ), bark ( $r = 0.682$ ,  $p < 0.05$ ) leaves (Y:  $r = 0.678$ ,  $p < 0.05$ ; O:  $r = 0.712$ ,  $p < 0.05$ ). Higher Fraction 2 resulted in higher plant uptake. Fraction 3 (oxidizable) had moderate correlation ( $r = 0.585$ ,  $p < 0.05$ ) and Fraction 4 (Residual) had high correlation with plant uptake in root ( $r = 0.749$ ,  $p < 0.05$ ) and bark ( $r = 0.776$ ,  $p < 0.05$ ). High correlation was found between young leaf and green leaf ( $r = 0.661$ ,  $p < 0.05$ ) and between green leaf and old leaf ( $r = 0.933$ ,  $p < 0.05$ ).

**Table 7.** Correlation between fractionation of trace metals in sediment and accumulation in plant parts

	Fraction1	Fraction2	Fraction3	Fraction4	Root	Bark	Leaf_Y	Leaf_M	Leaf_O
<b>Fraction1</b>	1								
<b>Fraction2</b>	-.590'								
<b>Fraction3</b>	-.711"	-.131							
<b>Fraction4</b>	-.228	.388	-.208						
<b>Root</b>	-.515	.634'	-.005	.749"					
<b>Bark</b>	-.480	.682'	-.089	.776"	.573				
<b>Leaf_Y</b>	-.798"	.458	.585'	.143	.313	.556			
<b>Leaf_M</b>	-.742"	.678'	.368	.031	.197	.506	.661'		
<b>Leaf_O</b>	-.627'	.712"	.201	.037	.249	.454	.470	.933"	1

Y = young leave, M = mature leave (green colour), O = old leave (yellow colour)

'Correlation is significant at the 0.05 level. "Correlation is significant at the 0.01 level

## Discussion

In this study, the samples were collected during the heavy rainy season and therefore several large freshwater bodies flowed into the lake and lessened the salinity of the lake water. However, the metal concentrations in the collected mangrove sediment samples in our study area as compared to the sediment quality guideline revealed that the As concentration was still higher than that of the quality guideline by 2-4 times and classified as heavily polluted. Comparison of our data regarding the metal concentrations (As, Cd, Pb and Zn) with the previous data from the other studies of Songkhla lake as well as other areas in the world was illustrated in *Table 8*. The maximum values of As and Pb at the Outer Section of Songkhla Lake previously reported by Pradit et al. (2010, 2013) were considerably lower than those of this study. Especially, As levels in Songkhla Lake were much higher than those found in Pattani bay (Pradit et al., 2016), Seitu Wetland (Pradit et al., 2016), Sundarban Wetland (Chowdhury et al., 2015) and at Kelantan Delta reported by Baruddin (2017). This result is certainly supported by Sompongchaiyakul and Sirinawin (2007), Pradit et al. (2010, 2013); who reported that concentrations of As, and Pb were ascending owing to rapid urban and industrial expansion. Outer Section of Songkhla Lake receives municipal wastes from two large and rapidly expanding cities of Songkhla and Hatyai as well as agricultural and industrial discharges transported by U-Taphao canal (Sompongchaiyakul and Sirinawin, 2007). The mean concentration values of As and Pb of this study were similar to the result of Pradit et al. (2010) but higher than those of Pradit et al. (2013). It is well accepted that the mangrove area is considered as a sink of trace metals and therefore the mangrove sediment samples of this study possesses the higher amounts of trace metals than those of the lake sediment reported by Pradit et al. (2013). Although the total concentration of metal is an appropriate indicator of contamination assessment, it does not provide enough information about bioavailability and toxicity of heavy metals (Zhong et al., 2011).

Chemical fractionation differentiates metals of natural origin from those derived from anthropogenic sources. The bioavailability fractions (exchangeable, oxidizable and reducible) of As, Cd and Pb in mangrove sediments of this study were greater than 90% whereas the residual fraction of these elements was very low. Thus, it was probably concluded that mangrove area in this study had high bioavailability. This may suggest that there was considerable anthropogenic input to the Songkhla Lake as compared to the low bioavailability of As (about 30%) reported by Baruddin et al. (2017). This was probably caused by different mangrove plant types. Our sampling area was *S. caseolaris* habitat whereas Baruddin et al. (2017) was *R. mucronata*.

In our study Pb seemed to be accumulated in plant parts. This was concordant with the study of Toledo-Bruno et al. (2016) in Mangrove forest reserve in Mindanao, Philippines discovering of *Sonneratia alba* being Pb-hyperaccumulator. Our study Pb was highest in root and well agreed with the study of Nazli and Hashim (2010).

Plants may be passive receptors of heavy metals, but they also exert control over uptake or rejection of some elements by appropriate physiological reactions. To illustrate this point, based on *Table 3*, the BAF values for all the studied heavy metals were always less than 1, which indicated that *Sonneratia caseolaris* exhibited restricted metal sediment-root uptake for non-essential heavy metals similar to BAF of *R. mucronata* in the Outer Section of Songkhla Lake (Baruddin et al., 2017)

**Table 8.** Trace metals concentrations ( $\mu\text{g/g}$  dry wt) in sediments of Songkhla Lake and at other sites

	As	Cd	Pb	Zn	References
Songkhla lake	20.4-50.5	0.006-0.182	48.8-78.8	48.6-126.6	This study
Songkhla Lake	5.1-25.7	-	-		Sompongchaiyakul and Sirinawin, 2007
Songkhla Lake	0.8 -70.7	0.1-2.4	8.2-131	5.4-561.6	Pradit et al., 2010
Songkhla Lake	20-22.0	0.21-0.28	31-35		Pradit et al., 2013
Songkhla Lake	20.7		55		Baruddin et al., 2017
Futian, China	152.4		70.7		Li et al., 2016
Pattani Bay, Thailand	2.98-9.34	0.00-0.02	2.40-11.48	0.01-7.06	Pradit et al., 2016
Seitu Wetland, Malaysia	0.89-3.52	0	0.63-1.56	0.50-2.30	Pradit et al., 2016
Kelantan Delta, Malaysia	18.28-34.95		31.30-75.84		Baruddin et al., 2017
Sundarban Wetland, India	3.22-4.41	0.19-0.22	11.6-20	32.51-36.33	Chowdhury et al., 2015
Background value	7	0.12	22.5	56	Choi et al., 2008
Heavily polluted	>8	>6	>60	>200	US EPA
Moderately polluted	3-8		<3	90-200	US EPA

## Conclusion

The anthropogenic inputs of As and Pb were high in the mangrove sediments of the lake especially at Kuannieng (station 2). As and Zn were found highest concentration in bark whereas Cd and Pb were found highest concentration in root. The BCF and TF values proposed that *Sonneratia caseolaris* showed limited (BCF < 1, and TF > 1) accumulation of As, Cd, Pb and Zn in their aboveground parts and thus presented a low food chain hazard except As and Cd at Phawong canal (station 3) and U-Taphao canal (station 4). The potential ecological risk index of single element indicated that As was classified as moderate risk level. The concentration of As should be concerned and regularly monitored.

**Acknowledgements.** This work was financially supported by the Prince of Songkla University under contract number COR590528S.

## REFERENCES

- [1] Baruddin, N. A., Shazili, N. A. M., Pradit, S. (2017): Sequential extraction analysis of heavy metals in relation to bioaccumulation in mangrove, *Rizophoramucronata* from Kelantan Delta, Malaysia. *Aquaculture, Aquarium, Conservation & Legislation*. – *International Journal of the Bioflux Society* 10: 172-181.
- [2] Baudo, R., Muntau, H. (1990): Lesser known in-place pollutants and diffuse source problems. – In: Baudo, R. Giesy, J. P., Muntau, H. (ed.) *Sediment Chemistry and Toxicity of In-Place Pollutants*, USA.
- [3] Choi, K. Y., Kim, S. H., Chon, H. T. (2008): Distributions and accumulations of heavy metals in the sediments of harbors and coastal areas in Korea. – *Proceedings of the*

- international symposia on geoscience resources and environments of Asian Terranes (GREAT 2008), 4<sup>th</sup> IGCP 516, and 5<sup>th</sup> APSEG. November 24-28, 2008. Bangkok, Thailand.
- [4] Chojnacka, K., Chojnacki, A., Gorecka, H., Gorecki, H. (2004): Bioavailability of heavy metals from polluted soils to plants. – *Science of the Total Environment* 337: 175-182.
- [5] Chowdhury, R., Favas, P. J., Pratas, J., Jonathan, M. P., Ganesh, P. S., Sarkar, S. K. (2015): Accumulation of trace metals by mangrove plants in Indian Sundarban Wetland: prospects for phytoremediation. – *International Journal of Phytoremediation* 17: 885-894.
- [6] Emsong Project (1998): Environmental diagnosis for the Songkhla lake basin: Technical Background Report No. 9. – VKI in association with: DHI, PEM consult A/S, COWI A/S, Prince of Songkla University and Seatec International Ltd.
- [7] Filgueiras, A. V., Lavilla, I., Bendicho, C. (2004): Evaluation of distribution, mobility and binding behaviour of heavy metals in surficial sediments of Louro River (Galicia, Spain) using chemometric analysis: a case study. – *Science of the Total Environment* 330: 115-129.
- [8] Håkanson, L. (1980): An ecological risk index for aquatic pollution control of sediment Ecological approach. – *Water Research* 14: 975-1000.
- [9] Hosani, M. A., Anouti, F. A. (2014): A Preliminary Exploration of Heavy Metal Contamination within *Avicennia marina* in the United Arab Emirates. – *Journal of Environmental & Analytical Toxicology* 4: 232. DOI: 10.4172/2161-0525.1000232.
- [10] Kim, R.-Y., Yoon, J.-K., Kim, T.-S., Yang, J. E., Owens, G., Kim, K.-R. (2015): Bioavailability of heavy metals in soils: definitions and practical implementation-a critical review. – *Environmental Geochemical Health* 37: 1041-1061.
- [11] Lanno, R., Wells, J., Conder, J., Bradham, K., Basta, N. (2004): The bioavailability of chemicals in soil for earthworm. – *Ecotoxicology and Environmental Safety* 57(1): 39-47.
- [12] Li, R., Chai, M., Qiu, G. Y. (2016): Distribution, fraction, and ecological assessment of heavy metals in sediment-plant system in mangrove forest, South China Sea. – *PloS ONE* 11(1). DOI: 10.1371/journal.pone.0147308.s.
- [13] Liu, J., Wu, H., Feng, J., Li, Z., Lin, G. (2014): Heavy metal contamination and ecological risk assessments in the sediments and zoobenthos of selected mangrove ecosystems, South China. – *CATENA* 119: 136-142.
- [14] Loring, D. H., Rantala, R. T. T. (1992): Manual for the geochemical analyses of marine sediments and suspended particulate matter. – *Earth-Science Reviews* 32: 235-283.
- [15] MacDonald, D. D., Ingersoll, C. G., Berger, T. A. (2000): Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. – *Archives of Environmental Contamination and Toxicology* 39: 20-31.
- [16] Macfarlane, G. R., Burchett, M. D. (2000): Cellular distribution of copper, lead and zinc in the grey mangrove, *Avicennia marina* (Forsk.) Vierh. – *Aquatic Botany* 68: 45-59.
- [17] Nath, B., Birch, G., Chaudhuri, P. (2013): Trace metal biogeochemistry in mangrove ecosystems: a comparative assessment of acidified (by acid sulfate soils) and non-acidified sites. – *Science of the Total Environment* 463: 667-674.
- [18] NaZli, M. F., Hashim, N. R. (2010): Heavy metal concentrations in an important mangrove species, *Sonneratia Caseolaris*, in Peninsular Malaysia. – *Environment Asia* 3: 50-55.
- [19] NOAA (1999): Sediment quality guidelines developed for the National Status and Trends Program. National Atmospheric Association. – [http://response.restoration.noaa.gov/book\\_shelf/121\\_sedi\\_qual\\_guide.pdf](http://response.restoration.noaa.gov/book_shelf/121_sedi_qual_guide.pdf).
- [20] Noriki, S. K., Nakanishi, T., Fukawa, M., Uematsu, T., Uchida, H., Tsunogai, S. (1980): Use of a teflon vessel for the decomposition followed by determination of chemical constituents of various marine samples. – *Bulletin of the Faculty of Fisheries Hokkaido University* 31: 345-465.
- [21] Pendias, A. K., Pendias, H. (2001): Trace metals in soils and plants. – CRC Press, Boca Raton.

- [22] Pradit, S., Gao, Y., Faiboon, A., Baeyens, W., Leermakers, M. (2013): Application of DET (diffusive equilibrium in thin films) and DGT (diffusive gradients in thin films) techniques in the study of the mobility of sediment-bound metals in the outer section of Songkhla Lake, Southern Thailand. – *Environmental Monitoring and Assessment* 185: 4207-4220.
- [23] Pradit, S., Sangmanee, W., Towatana, P., Shazili, N. A. M. (2016): Trace metals, grain size and organic matter in sediment from a coastal area in Pattani Bay, Thailand and setiuwetland, Malaysia. – *Aquatic Ecosystem Health and Management* 19: 345-354.
- [24] Pradit, S., Wattayakorn, G., Angsupanich, S., Baeyens, W., Leermakers, M. (2010): Distribution of trace elements in sediments and biota of Songkhla Lake, Southern Thailand. – *Water, Air, and Soil Pollution* 206: 155-174.
- [25] Rauret, G., López-Sánchez, J. F., Sahuquillo, A., Rubio, R., Davidson, C., Ure, A., Quevauviller, P. (1999): Improvement of the BCR three-step sequential extraction procedure prior to the certification of new sediment and soil reference materials. – *Journal of Environmental Monitoring* 1: 57-61.
- [26] Sarkar, S. K., Favas, P. J. C., Rakshit, D., Satpathy, K. K. (2014): Geochemical Speciation and Risk Assessment of Heavy Metals in Soils and Sediments. – In: Maria, C., Hernandez, S. (eds.) *Environmental Risk Assessment of Soil Contamination*. InTech Publishing, Rijeka.
- [27] Sekomo, C. B., Nkuranga, E., Rousseau, D. P. L., Lens, P. N. L. (2011): Fate of heavy metals in and urban natural wetland: The Nyabugogo Swamp (Rwanda). – *Water, Air, & Soil Pollution* 214: 321-333.
- [28] Shazili, N. A. M., Mohamed, K. A. R., Husain, M. L., Asmawi, N., Salmah, A. (1999): Tracemetals in the surface sediments of the South China Sea, Area I: Gulf of Thailand and the East Coast of Peninsular Malaysia. – In: *Proceedings of the First Technical Seminar on Marine Fishery Resources Survey in the South China Sea Area I Gulf of Thailand and East Coast of Peninsular Malaysia*, pp. 73-85. Southeast Asian Fisheries Development Center, Samutprakan, Thailand.
- [29] Sojisuporn, P. (2005): Sediment and shallower problems. In the master plan for Songkhla Lake basin development. – *Water Resources* 5: 167-191.
- [30] Sompongchaiyakul, P., Sirinawin, W. (2007): Arsenic, Chromium and Mercury in surface sediment of Songkhla Lake system, Thailand. – *Asian Journal of Water, Environment and Pollution* 4(1): 17-24.
- [31] Takac, P., Szabova, T., Benkova, M. (2009): Heavy metals and their bioavailability from soils in the long-term polluted central spis region of SR. – *Plants Soil Environment* 55(4): 167-172.
- [32] Tokalioglu, S., Senol, K., Latif, E. (2000): Determination of heavy metals and their speciation in lake sediments by Flame Atomic Absorption Spectrometry after a four-stage sequential extraction procedure. – *AnalyticaChimicaActa* 413: 33-40.
- [33] Toledo-Bruno, A. G. I., Aribal, L. G., Lustria, M. G. M., Marin, R. A. (2016): Phytoremediation potential of mangrove species at Pangasihang Mangrove forest reserve in Mandanao, Philippines. – *Journal of Biodiversity and Environmental Sciences* 9: 142-149.
- [34] USEPA (1989): *Assessing Human Health Risks from Chemically Contaminated Fish and Shellfish. A Guidance Manual*. – EPA503/8-89-002. USEPA, Washington, D. C.
- [35] Zhong, X. L., S L Zhou, Q. Zhu, Zhao, Q. G. (2011). Fraction distribution and bioavailability of soil heavy metals in the Yangtze River Delta-A case study of Kushan City in Jiangsu Province, China. – *Journal of Hazardous Materials* 198: 13-21.
- [36] Zhou, Y. W., Zhao, B., Peng, Y. S., Chen, G. Z. (2010): Influence of mangrove reforestation on heavy metal accumulation and speciation in intertidal sediments. – *Marine Pollution Bulletin* 60: 1319-1324.