

EFFECT OF NITROGEN FERTILIZATION LEVELS ON GRAIN YIELD AND YIELD COMPONENTS IN TRITICALE BASED ON AMMI AND GGE BILOT ANALYSIS

ORAL, E.

*Department of Seed Production, Kızıltepe Vocational Training High School, Artuklu University
21500 Kızıltepe, Mardin, Turkey
(e-mail: eroloral65@gmail.com; phone: +90-546-746-6266)*

(Received 22nd May 2018; accepted 17th Jul 2018)

Abstract. This study examined the effects of nitrogen doses rate on grain yield (GY) and yield components of spring triticale (*×Triticum secale*) cultivars in dry area, based on cultivars plus cultivar × nitrogen doses interaction GGE biplot and AMMI analysis. The research was designed to evaluate the effects of different nitrogen fertilization levels (0, 40, 80, 120 and 160 kg ha⁻¹ of N) on the grain yield and agronomic performance of two cultivars in two growing seasons (2015-16/2016-17). Split plot layout with in randomized complete block design with 3 replications was used in both years. The response to fertilization levels were evaluated through GGE (Genotype main effects and Genotype x Environment interaction) biplot graphic methodologies and regression. Combined analysis of variance of nitrogen applications of two cultivars showed highly significant ($p < 0.01$) difference between the cultivars, nitrogen applications and interaction. The grain yield of triticale was most of all affected by cultivars (70.8%) and then by nitrogen doses (23.5%) and interaction (5.7%). On the other hand, the study showed that plant height, number of spikes per m², number of grain of ears and grain yield increased depending on the increasing nitrogen doses of the cultivars. The highest performance of grain yield of both cultivars were associated with nitrogen N₃ (120 kg/ha) fertilization in regression and AMMI analysis. The results showed that 80 kg/ha N application can be recommended in triticale traits and grain yield with Esin (new) cultivar. Further more, we found that the GGE biplot method generated highly useful results with high visual quality in the study.

Keywords: *triticale, nitrogen doses, yield, yield components*

Introduction

Triticale is a new species in field crops and it is expected to be regarded as a seed in the future as well as being evaluated as forage. It is already produced for seed purposes in many African countries. Especially in dry farming areas, where barley and wheat are not well cultivated, in areas where extreme temperatures are experienced and rainfall is insufficient, triticale can be more economically grown than other cereal crops. In these places, triticale is contribute the several advantages in cropping system because cereal grains can use N, captured in soil profile after lentil or chickpea crops. After lentil and chickpea farming it can use N to accumulate in the soil. In addition, because of the fact that the triticale is abstinent plant, so it is able to get better nitrogen from soil in dry areas compared to wheat and barley. The Syrian border, where the research was carried out, is quite dry and it has become almost impossible to grow wheat without watering. Under these conditions, farmers tend to other cereal crops that may be an alternative to wheat in arid conditions. For this reason, growing triticale is becoming increasingly important (Kendal, 2015; Kendal et al., 2016).

Triticale was designed in order to both biotic and abiotic stressors and thus more suitable for cultivation in marginal areas than wheat (Kendal et al., 2012; Villegas et al., 2010). Triticale is often reported as an interesting product for adverse environmental conditions where environmental factors limit product cultivation (Ugarte et al., 2007;

Estrada-Campuzano et al., 2008). In such places, each plant does not use the same nitrogen in the same way and the plant cannot make maximum use application amount of nitrogen. Proper N management, are important for obtaining optimum crop yields and positive economic returns while limiting negative environmental impacts of crop production (Gibson et al., 2007). Many authors have reported positive effect of N fertilization on grain yield and the studies showed that N fertilization is one of the most important crop management techniques for spring triticale production (Gülmezoglu, Kinaci, 2004; Lestingi et al., 2010). The weather conditions is affect spring triticale grain yield performance and nitrogen (N) fertilization effect total 35.7% grain yield increase, compared with unfertilized spring triticale (Janušauskaitė, 2013).

Nitrogen is needed for early tiller development of plant to set up the crop for a high yield potential. On the other hand; nitrogen fertilization has an important effect on the final harvest, thus if this element is not take from plant, yield is decreases (Moreno et al., 2003). The amount of nitrogen, barley crop needs to reach maximize yield and quality, will depend on the seasonal conditions, soil type, and rotational history of the soil as well as the potential yield of the cultivars. Nitrogen is the key nutrient input for achieving higher yield of triticale (Alazmani, 2015). The farmers have not enough information about use nitrogen fertilizers and adequate information concerning actual soil requirements. Therefore, the study of use N dozes in triticale cultivars is necessary to recommend optimum nitrogen doses for high yield and yield components in different environment conditions.

The yield of each variety in any environment is a sum of environment (E) main effect, genotype (G) main effect and genotype by environment interaction (GE or GEI) (Farshadfar et al., 2013; Kilic, 2014; Sayar et al., 2013). On the other hand; farmers need varieties that show high performance in terms of yield and other essential agronomic traits by use nitrogen fertilizer. Different statistical analysis, such as correlation, path coefficient and principal component analysis (PCA) can be used to reveal associations between yield and other agronomic traits. The impact of GGE Biplot methods and regression analysis has been clearly showed by different researchers using relationship among factors. This methods; provide the correlative size and interaction (Asfaw et al., 2009; Sayar and Han, 2015; Kendal and Sayar, 2016). So it is very important to identify the use of nitrogen fertilization doses to cultivars for high yield and best quality. The objective of study is to determine the effect of different N rates on yield, yield components, and to identify optimal N deses rate using GGE Biplot analysis to recommend doses for application in Southeast Anatolia Region of Turkey.

Material and methods

The experiment was conducted in the research field of the Kiziltepe Vocational and Training High School, Mardin, Turkey. The experiment was conducted on the basis of split plot layout with completely randomized block design with 3 replications. Main plot was different level of nitrogen fertilizer (0, 40, 80, 120 and 160 kg ha⁻¹ of N and sub plot was different two spring triticale cultivars (*Table 1*).

This research was conducted in 2015-2016 and 2016-2017 growing seasons. The seeding rates were 500 seeds m⁻². Plot size was 7.2 m² (1.2 × 6 m) consisting of 6 rows spaced 20 cm apart. Sowings were made by using an experimental drill. The fertilization rates for all plots were different N ha⁻¹ doses and 60 kg P ha⁻¹ with sowing time and different N ha⁻¹ doses was applied to plots in double ridge stage. Harvests were

made using Hege 140 harvester in 6 m². Other normal agronomic practices for barley production were followed. During both of growing seasons, heading time (date), spike per square (m²), maturity time (date) plant height (cm), length of spike (cm), number of grains per spike, thousand grain weight (g), grain yield (kg/ha⁻¹), hectoliter weight (kg hl), Protein Content (%) were examined (Kendal and Sayar, 2016). The soil analysis results are shown in *Table 2* and the climate data of growing seasons are shown in *Table 3* (Mardin Regional Directorate of Meteorology). Different nitrogen use efficiency (NUE, kg kg⁻¹ N) was calculated for each treatment: $NUEX\ Trait = (X\ trait\ N - X\ trait\ N0) / N_x$, where X trait N is X trait from nitrogen (N) fertilized treatments, Xtrait N0 – X trait in unfertilized treatment, and N_x– nitrogen input (N40, N80, N120, N160, kg ha⁻¹ N).

Table 1. Information about cultivars used in the experiment

Cult. name	Where registration	Regist. time	Type
Tacettinbey	Cukurova University	2000	Spring
Esin	GAP International Agricultural and Training Center	2016	Spring

Table 2. Soil analysis results of trial

Texturing (%)			pH	Organic subs. (%)	Total P ppm	Total K ppm	Lime%
Sand	Clay	Mile					
65.6	22.4	12.0	7.5	1.4	22	210	17.15

Table 3. Information about meteorological data of location of study

Months	Humidity (%)			Temperature (°C)			Precipitation (mm)		
	2015-16	2016-17	LTA*	2015-16	2016-17	LTA	2015-16	2016-17	LTA
October	49.6	33.3	32.9	19.5	20.5	19.3	58.2	16.5	19.7
November	50.3	35.2	50.2	11.7	11.5	11.2	62.0	27.2	49.1
December	51.7	71.3	47.2	6.4	3.2	4.2	64.7	128.4	58.5
January	74.1	63.6	63.7	2.2	3.1	3.2	146.3	38.3	78.7
February	66.2	51.3	51.0	8.5	3.8	3.9	3.6	23.2	64.4
March	59.1	62.5	62.9	10.0	9.7	9.0	119.8	101.7	99.6
April	41.3	55.7	55.2	16.8	13.5	15.2	27.1	109.2	98.5
May	42.0	44.0	43.8	19.8	19.7	19.6	20.0	60.3	57.0
June	28.2	26.1	25.8	26.2	26.8	26.0	1.0	0.2	2.2
July	22.4	17.0	16.5	30.6	32.4	32.1	0.1	0.0	0.6
Total							502.8	505.0	528.3
Mean	48.49	46.00	44.9	15.7	14.4	14.3			

LTA: long term average, *Mardin regional directorate of meteorology station (2017)

The data obtained from the study related to the investigated grain yield and yield components were analyzed respectively for each year and combined with nitrogen doses by using the JMP 5.0.1 statistical software package (SAS Institute, 2002), and the differences between means were compared using a least significant difference (LSD) test at the 0.05 probability level (Steel and Torrie, 1980). Also regression analysis was done by this program. On the other hand; GGE biplot analyses were carried out using GGE biplot software to show the differences among the applied nitrogen doses and crop characteristics and cultivar crop characteristics in two growing seasons (Dogan et al., 2016; Kilic, 2016). GGE biplot analysis also allows comparison amongst nitrogen doses in terms of their discriminating ability and representativeness. These values can be assessed using the discriminating power of the doses' biplot screen of the GGE biplot (Yan and Thinker, 2006). With the AMMI biplot analysis graph in the study: It was aimed at illustrating grain yield performance of N doses and status of triticale cultivars (*Fig. 1b*), grain yield performance of cultivar (*Fig. 1c*), the effect of N doses on grain yield on years (*Fig. 1d*). Also, with the GGE biplot analysis graphs in the study: It was aimed at revealing relation among examined doses and traits (*Fig. 1e*), sector analysis of doses and grouping traits (*Fig. 1f*), ranking of doses on traits (*Fig. 1g*), comparison of doses on traits (*Fig. 1h*)

Results

The combined ANOVA revealed highly significant differences among the years, cultivars and interaction of them for all components ($P < 0.01, 0.05$), the differences among nitrogen doses were highly significant ($P < 0.01, 0.05$) for all components as shown in *Tables 4* and *5*. Generally, breeders interested in the genotypes with high genotypic main effect (average over years and nitrogen doses) and with low fluctuation in yield or yield components (stable). On the other hand, the study showed that plant height, number of spikes per m^2 , number of grain of ears and grain yield increased depending on the increasing nitrogen doses of the cultivars.

The results of the data reviewed

Table 4. Variance of analysis on grain yield and yield components of triticale

Sources	DF	HT (date)	SS (m^2)	MT (date)	PH (cm)	SL (cm)	NGS	TGW (g)	GY (kg/ha^{-1})	HL (kg/hl^1)	PC (%)
Year	1	400.4**	1696.0*	589.1**	779.0*	3.3ns	477.7*	0.9ns	522.2ns	123.8*	0.3ns
Error 1	4	0.7	413.9	0.9	36.7	3.5	14.4	9.1	4322.9	4.4	3.6
Cultivar	1	88.8**	380.0ns	26.7**	449.4**	17.1**	492.5**	212.8**	158620.0**	13.1*	1.5*
Year*Cult.	1	4.8**	498.8ns	6.7**	2.5ns	2.4ns	273.5**	28.8**	84.0ns	86.9**	1.3*
Nitr. Doses	4	2.4*	7083.2**	2.5*	2607.4**	58.9**	391.5**	20.4*	52602.6**	33.4*	5.9*
Year*N.Dos	4	1.2ns	5371.2*	1.1*	132.1*	11.6*	69.5*	3.2ns	8718.4**	18.2*	0.5ns
Cul.*N.Dos	4	0.4ns	130.9ns	0.5 ns	57.9ns	10.8ns	47.3ns	40.3**	12757.2**	12.5*	1.1ns
Y*C*ND	4	0.1ns	1247.4ns	0.8ns	120.6*	2.8ns	18.7ns	10.8ns	410.9ns	19.5*	1.2ns
Error 2	36	5.9	6039.5	3.7	267.8	23.1	143.8	49.6	6022.4	40.5	7.5
C. Total	59	504.9	22861.0	632.0	4453.4	133.4	1928.9	376.0	244061.0	352.2	23.0
CV(%)		0.24	9.18	1.03	2.52	7.22	4.37	3.14	3.36	1.37	4.22

HT: Heading Time, SS: Spike ofper Square, MT: Maturity time, PH: Plant Height, LS: Length of Spike, NGS: Number of Grains per Spike, TGW: Thousand Grain Weight, GY: Grain Yield, HL: Hectoliter weight, PC: Protein Content

Table 5. The variance of AMMI analysis on grain yield of nitrogen doses

Source	Df	SS	MS	F	Explained SS (%)
Total	59	244061	4137		
Treatments	9	223980	24887	70.23	
Cultivar	1	158620	158620	447.65**	70.8
N doses	4	52603	13151	22.26**	23.5
Block	10	5907	591	1.67	
Interactions	4	12757	3189	9.00**	5.7
IPCA1	4	12757	3189	9.00**	
Error	40	14174	354		

Df: degrees of freedom; **: $p < 0.01$; *: $P < 0.05$

The mean grain yield (GY) of 2015/16 growing season (3878 kg ha^{-1}) was higher than that of 2016/17 (3819 kg ha^{-1}), the mean yield of growing seasons was changed from 3334 kg ha^{-1} (Esin) to 4362 kg ha^{-1} (Tacettinbey) in Table 6.

Table 6. The data effect interaction of nitrogen levels and triticale cultivars

Cultivar		N Doses					Mean		N Doses					Mean
		N ₀	N ₁	N ₂	N ₃	N ₄			N ₀	N ₁	N ₂	N ₃	N ₄	
Tacettinbey	HT (date)	163.5	163.8	163.8	163.8	164.2	163.8 ^B	NGS	39.0 ^d	40.9 ^{cd}	42.6 ^c	46.0 ^b	46.0 ^b	42.8 ^B
Esin		166.0	166.0	166.5	166.3	166.5	166.3 ^A		43.2 ^c	49.5 ^a	50.2 ^a	51.0 ^a	48.8 ^a	48.6 ^A
Mean		164.8 ^C	164.9 ^{BC}	165.2 ^{AB}	165.1 ^{AC}	165.3 ^A			41.1 ^D	45.2 ^C	46.4 ^{BC}	48.5 ^A	47.4 ^{AB}	
Tacettinbey	SS (m ²)	125.8	134.0	140.2	137.7	154.8	138.5	TGW	34.9 ^{cd}	35.9 ^{cd}	36.0 ^c	35.8 ^{cd}	34.5 ^d	35.4 ^B
Esin		128.7	136.7	144.0	142.8	165.5	143.5		36.4 ^b	37.2 ^b	37.1 ^b	38.0 ^a	38.2 ^a	39.2 ^A
Mean		127.2 ^C	135.3 ^{BC}	142.1 ^B	140.3 ^B	160.2 ^A			36.4 ^C	36.9 ^{BC}	37.3 ^{AC}	38.0 ^A	37.8 ^{AB}	
Tacettinbey	MT(date)	30.0	30.2	30.3	30.5	30.7	31.7 ^A	GY(kg/ha)	2948 ^h	3542 ^{ef}	3408 ^{fg}	3485 ^f	3287 ^g	3334 ^B
Esin		31.5	31.5	31.8	31.5	32.0	30.3 ^B		364.8 ^e	4200 ^d	4657 ^b	4847 ^a	4460 ^c	4362 ^A
Mean		30.8 ^C	30.8 ^{BC}	31.1 ^{AB}	31.0 ^{BC}	31.3 ^A			329.8 ^D	3871 ^C	4033 ^B	4166 ^A	3873 ^C	
Tacettinbey	PH (cm)	95.9	100.5	103.5	112.5	114.7	105.4 ^B	HW	74.3 ^b	76.9 ^a	76.9 ^a	77.2 ^a	77.6 ^a	76.6 ^B
Esin		102.1	105.6	112.0	117.5	117.2	110.9 ^A		77.0 ^a	77.6 ^a	77.4 ^a	78.0 ^a	77.7 ^a	77.5 ^A
Mean		99.0 ^D	103.1 ^C	107.7 ^B	115.0 ^A	116.0 ^A			75.6 ^B	77.3 ^A	77.2 ^A	77.6 ^A	77.6 ^A	
Tacettinbey	LS (cm)	8.3 ^f	10.2 ^e	10.7 ^{de}	12.3 ^{ab}	11.3 ^{cd}	10.6 ^B	PC (%)	10.5	11.1	10.8	11.5	11.1	11.0 ^A
Esin		10.3 ^e	11.5 ^{bd}	12.5 ^a	12.2 ^{bc}	11.7 ^{bc}	11.6 ^A		10.2	10.3	10.8	11.0	11.0	10.7 ^B
Mean		9.3 ^D	10.8 ^C	11.6 ^B	12.3 ^A	11.5 ^B			10.3 ^C	10.7 ^{BC}	10.8 ^B	11.3 ^A	11.0 ^{AB}	

HT: Heading Time, SS: Spike per Square, MT: Maturity time, PH: Plant Height, LS: Length of Spike, N Doses – N₀: not applied, N₁: 40 kg/ha, N₂: 80kg/ha, N₃: 120 kg/ha N₄: 160 kg/ha

The yield of application nitrogen doses were ranged from 3298 kg ha^{-1} (N₀) to 4166 kg ha^{-1} (N₃). The yield of cultivar and nitrogen interaction were changed from 2948 kg ha^{-1} to 4847 kg ha^{-1} and the best yield was obtained by N₃ nitrogen doses in Esin, while the low yield obtained from N₀ (without nitrogen) doses and Tacettinbey variety (Table 7).

Table 7. Influence of different nitrogen levels on yield and yield components of triticale cultivars

Cultivar		2015-2016					Mean	2016-2017					Mean
		N0	N1	N2	N3	N4		N0	N1	N2	N3	N4	
Tacetinbey	HT (date)	161.0	161.3	161.7	161.7	162.0	161.5 ^D	166	166.3	166	166	166.3	166.1 ^B
Esin		163.0	163.0	163.7	163.7	163.7	163.4 ^C	169	169	169.3	169	169.3	169.1 ^A
N mean		162.0 ^c	162.2 ^c	162.7 ^b	162.7 ^b	162.8 ^b		167.5 ^a	167.7 ^a	167.7 ^a	167.5 ^a	167.8 ^a	
Year mean		162.5 ^B						167.6 ^A					
Tacetinbey	SS (m ²)	121.0	124.7	134.0	139.3	161.3	136.1	130.7	143.3	146.3	136	148.3	140.9
Esin		107.0	110.3	138.7	147.0	173.7	135.3	150.3	163	149.3	138.7	157.3	151.7
N mean		114.0 ^d	117.5 ^d	136.3 ^c	143.2 ^{bc}	167.5 ^a		140.5 ^{bc}	153.2 ^{ab}	147.8 ^{bc}	137.3 ^c	152.8 ^{ab}	
Year mean		135.7 ^B						146.3 ^A					
Tacetinbey	MT (date)	27.0	27.0	27.7	28.0	28.0	27.5 ^D	33.0	33.3	33.0	33.0	33.33	33.1 ^B
Esin		28.0	28.0	28.3	28.0	28.7	28.2 ^C	35.0	35.0	35.3	35.0	35.33	35.1 ^A
N mean		27.5 ^c	27.5 ^c	28.0 ^b	28.0 ^b	28.3 ^b		34.0 ^a	34.2 ^a	34.2 ^a	34.0 ^a	34.3 ^a	
Year mean		27.9 ^B						34.1 ^A					
Tacetinbey	PH (cm)	92.1 ⁱ	95.3 ^{hi}	96.9 ^h	111.6 ^{df}	112.1 ^{de}	101.6	99.7 ^h	105.7 ^g	110.0 ^{eg}	113.3 ^{ce}	117.3 ^{bd}	109.2
Esin		96.9 ^h	99.3 ^h	110.9 ^{ef}	114.3 ^{be}	116.0 ^{bd}	107.5	107.3 ^{fg}	112.0 ^{de}	113.0 ^{ce}	120.7 ^a	118.3 ^{ab}	114.27
N mean		94.5 ^F	97.3 ^F	103.9 ^E	112.9 ^C	114.1 ^{BC}		103.5 ^E	108.8 ^D	111.5 ^{CD}	117.7 ^{AB}	117.8 ^A	
Year mean		104.5 ^B						111.7 ^A					
Tacetinbey	SL (cm)	8.0	9.3	9.7	12.3	11.3	10.1	8.7	11.0	11.7	12.3	11.3	11.0
Esin		9.7	11.0	12.3	13.0	12.0	11.6	11.0	12.0	12.7	11.3	11.3	11.7
N mean		8.8 ^F	10.2 ^{DE}	11.0 ^{CD}	12.7 ^A	11.7 ^{BC}		9.8 ^E	11.5 ^{BC}	12.2 ^{AB}	11.8 ^{AC}	11.3 ^{BC}	
Year mean		10.9						11.3					
Tacetinbey	NGS	38.6	41.3	41.5	44.6	44.8	42.2 ^B	40.3	44.7	45.0	44.0	43.7	43.5 ^B
Esin		39.4	40.5	43.7	47.3	47.2	43.6 ^B	46.0	54.3	55.3	58.0	54.0	53.5 ^A
N mean		39.5 ^e	43.0 ^d	43.2 ^d	44.3 ^d	44.3 ^d		42.7 ^d	47.4 ^c	49.5 ^{bc}	52.7 ^a	50.6 ^{ab}	
Year mean		43.0 ^B						48.5 ^A					
Tacetinbey	TGW (1000)	36.1	36.6	36.9	35.9	34.3	36.0 ^C	33.6	35.1	35.1	35.6	34.7	34.8 ^D
Esin		36.6	36.5	38.3	39.8	40.5	38.3 ^B	39.2	39.3	39.1	40.5	41.8	40.0 ^A
N mean		36.4	36.6	37.6	37.9	37.4		36.4	37.2	37.1	38.0	38.2	
Year mean		37.2						37.4					
Tacetinbey	GY (kg/ha)	3087	3773	3403	3410	3203	3375 ^B	2810	3310	3413	3560	3370	3293 ^B
Esin		3667	4437	4687	4760	4350	4380 ^A	3630	3963	4627	4933	4570	4345 ^A
N mean		3377 ^D	4105 ^{AB}	4045 ^B	4085 ^B	3777 ^C		3220 ^E	3637 ^C	4020 ^B	4247 ^A	3970 ^B	
Year mean		3878						3819					
Tacetinbey	HW (kg/hl)	75.4 ^{df}	76.7 ^{ce}	76.8 ^{ce}	75.2 ^{ef}	77.6 ^{be}	76.3 ^B	73.1 ^g	77.1 ^{cd}	77.1 ^{cd}	79.3 ^{cd}	77.5 ^{ab}	76.8 ^B
Esin		74.3 ^{fg}	74.5 ^{fg}	74.3 ^{fg}	75.5 ^{df}	75.7 ^{df}	74.9 ^C	79.6 ^a	80.7 ^a	80.4 ^a	80.5 ^a	79.6 ^a	80.2 ^A
N mean		74.9 ^E	75.6 ^{CE}	75.6 ^{CE}	75.3 ^{DE}	76.7 ^C		76.3 ^{CD}	78.9 ^{AB}	78.8 ^{AB}	79.9 ^A	78.6 ^B	
Year mean		75.6 ^B						78.5 ^A					
Tacetinbey	PC (%)	10.2	10.8	10.6	11.4	10.9	10.8 ^B	10.7	11.3	11.0	11.6	11.4	11.2 ^A
Esin		10.6	10.5	10.6	10.9	11.1	10.7 ^B	9.8	10.1	11.1	11.1	10.8	10.6 ^B
N mean		10.4	10.7	10.6	11.1	11.0		10.3	10.7	11.0	11.4	11.1	
Year mean		10.8						10.9					

NGS: Number of Grains per Spike, TGW: Thousand Grain Weight, GY: Grain Yield, HL: Hectoliter Weight, PC: Protein Content. N Doses – N₀: not applied, N₁: 40 kg/ha, N₂: 80 kg/ha, N₃: 120 kg/ha, N₄: 160 kg/ha

The results of grain yield showed that first growing season had high yield than the second year and Esin variety was yielding than Tacettinbey during two growing seasons. The results of grain yield showed that first growing season had higher yield than the second year and Esin variety changed from 4760 kg/hl to 4933 kg/hl (N₃-2015/16-2016/17) had a higher yield Tacettinbey (2810-3087 kg/hl, N₀) during two growing seasons in *Table 7*. Nitrogen use efficiency (NUE) was considerably influenced by cultivars, using N doses and growing seasons. The main factor determining the differences in NUE yield of cultivars was changed from 13.1% (Tacettinbey) to 19.6% (Esin) (*Table 7*). The influence of growing season on NUE_{yield} was changed from 21.6% (2015-16) to 23.3% (2016-17). The highest NUE_{yield} of interaction was obtained from N₃ fertilization rate (35.9%) in 2016-17 growing season. As expected, depend on increasing basic N fertilization rate increased NUE_{yield}. Each N rate increasing by 40 kg increased NUE_{yield} by 2.5 kg kg⁻¹.

The mean thousand grain weight (TGW) of 2016/17 growing season (37.4 g) was high than 2015/16 (37.2 g), the mean thousand grain weight of cultivars of both years was changed from 34.8 g (Tacettinbey) to 40.0 g (Esin) in *Table 6*. The number of thousand grain weight of nitrogen doses and cultivar interaction of both year means were changed from 36.4 g (N₀) to 38.2 g (N₄, 2016/17). The application nitrogen doses of thousand grain weight were ranged from 36.4 g (N₀) to 38.0 g (N₃) in *Table 7*. The thousand grain weight was increased depend on nitrogen rate till N₃ application doses for both varieties.

The mean hectoliter weight (HW) of 2016/17 growing season (78.5 kg/hl) was high than 2015/16 (75.6 kg/hl), the mean hectoliter weight of cultivars of both years was changed from 76.8 kg/hl (Tacettinbey) to 80.2 kg/hl (Esin) in *Table 6*. The number of hectoliter weight of nitrogen doses and cultivar interaction of both year means were changed from 74.9 kg/hl (N₀-2015/16) to 79.9 g (N₃-2016/17). The application nitrogen doses of hectoliter weight were ranged from 75.6 kg/hl (N₀) to 77.6 kg/hl (N₃ and N₄) in *Table 7*. The hectoliter weight was increased depend on nitrogen rate till N₄ doses for both varieties. N₃ and N₄ doses had the same values, so the N₃ dose application is better for hectoliter weight in triticale.

The mean protein content (PC) of 2016/17 growing season (10.9%) was higher than 2015/16 (10.8%), the mean hectoliter weight of cultivars of both years was changed from 11.2% (Tacettinbey) to 10.6% (Esin) in *Table 6*. The number of protein content of nitrogen doses and cultivar interaction of both year means were changed from 10.3% (N₀-2016/17) to 11.4% (N₃-2016/17). The application nitrogen doses of protein content were ranged from 10.3% (N₀) to 11.3% (N₃) in *Table 7*. The protein content was increased depend on nitrogen rate till N₃ doses for both varieties. Nitrogen use efficiency (NUE) of protein content influenced by cultivars, using N doses and growing seasons. The main factor determining the differences in NUE_{protein} of cultivars was changed from 5.3% (Tacettinbey) to 4.2% (Esin) (*Table 7*). The influence of growing season on NUE_{protein} was changed from 4% (2015-16) to 7% (2016-17). The highest NUE_{protein} of interaction was obtained from N₃ fertilization rate (10.7%) in 2016-17 growing season. As expected, depend on increasing basic N fertilization rate increased NUE_{protein}. Each N rate increasing by 40 kg increased NUE_{protein} by 1.5-2.5 kg kg⁻¹ till N₃ fertilization rate

The heading time (HT) was calculated from 01 January to heading time as day for both varieties separately in *Table 7*. The maturity time (MT) was calculated from heading time to maturity time as day for both varieties separately. On the other hand,

ear per square (SS), plant height (PH), long of spike (LE), number of grain per spike (NGS) were recorded and results were showed in *Table 7*. The results showed that there is positive correlation between heading time and maturity time, and depend on N application doses heading time and maturity time are delayed. The cultivars are affected the same on application doses. On the other hand the study indicated the ear per square (SS) and grain per spike (NGS) are contributing the grain yield. Moreover, all traits results (HT, MT, LS, NGS and SS) were increased depend on nitrogen rate till N₃ doses for varieties, years and interaction.


AMMI and GGE biplot analysis

Analysis of variance for nitrogen doses (ND) component (C), and the cultivar (C) × component (C) interaction showed significant ($P < 0.01$, $P < 0.01$) effect, and the total sum of squares explained for 96.15%, with PC1 and PC2 accounting 85.03% and 11.12% for nitrogen doses (ND) component (C) (*Fig. 1e-h*). Moreover, the AMMI analysis showed that the grain yield of triticale was most of all affected by cultivars (70.8%) and then by nitrogen doses (23.5%) and interaction (5.7%).


AMMI Analysis was done on grain yield for interaction nitrogen doses, cultivars and years (*Fig. 1b-e*). The results of interaction nitrogen doses and cultivars of grain yield (*Fig. 1b*) showed that grain yield is increased depend on nitrogen doses (*Fig. 1a*) till N₃ application doses, the contribute of N₁ and N₄ to grain yield of triticale is the same, but N₃ application dose is contribute to grain yield in highest level. The interaction of N doses and years showed that the highest grain yield was obtained from the first year (*Fig. 1c*), and the effect of N₃ doses on grain yield was highest than other N doses. The interaction of cultivars and years on grain yield indicated that first year (2015/16) is yielding than the second year (2016/17) and Esin variety is yielding than Tacettinbey variety for both years (*Fig. 1d*).

GGE biplot analysis shows the means over years for nitrogen doses relationships among yield components (*Fig. 1e-h*), the results of relationship between N doses and yield components showed that there is high correlation among components and N application doses (*Fig. 1e*). The figures show that there is high correlation between N₂ application dose and all yield components, while there is high correlation between N₃ doses and GY, LS, NGS, HW TGW and high correlation between N₄ and MT, PC, HT, PH and there is not any correlation among N₀, N₁ and yield components. The sector analysis of N doses and grouping of traits showed in *Fig. 1f*, the results indicated that four sectors occurred on the figure and MT, PC, HT, PH took places in first group in first sector with N₄, and GY, LS, NGS, HW TGW took places in second group in second sector with N₃, while N₂ took places in center of first and second sector, so this dose took places on center of all components. On the other hand N₀ and N₁ located in different sector and opposite of component group, so did not related with any component. The ranking biplot of N doses on components showed in *Fig. 1g*, and the figure indicated that N₂ dose is stable for all components, because N₂ dose located center of stable line with locate above mean of components line, while N₂ doses located side of GY, LS, NGS, HW TGW and desirable for these components, and N₂ desirable for MT, PC, HT, PH. On the other hand, N₀ and N₁ located in under mean of component line, so these doses are undesirable for all components. The comparison biplot of N doses on components showed in *Fig. 1h*, and the figure indicated that N₂ dose is stable for all components, because N₂ dose located on ideal enter line, while N₃ doses located side of GY, LS, NGS, HW TGW and desirable for these components, and N₄ desirable


for MT, PC, HT, PH. On the other hand, N₀ and N₁ located in under mean of component line, so these doses are undesirable for all components.




a




b




c



d



e



f




Figure 1. *a.* Regression analysis of grain yield (kg/ha) of cultivar and nitrogen doses. *b.* AMMI analysis of N doses and cultivars on grain yield. *c.* AMMI analysis on grain yield of cultivar. *d.* AMMI analysis of N doses of years. *e.* The relation between N doses and traits. *f.* The sector analysis of N doses and grouping of traits. *g.* Ranking of N doses on traits. *h.* Comparison of N doses on traits

Discussion

Analysis of variance for grain yield and yield components showed that nitrogen dose, year and cultivar was the main independent factor determining the differences in values of parameters between N doses and results. The significant differences ($P < 0.001$, $P < 0.005$) found of the yield and yield components and the grain yield of triticale was most of all affected by cultivars (70.8%) and then by nitrogen doses (23.5%) and interaction (5.7%) in *Table 5*, therefore the effect of nitrogen doses gives a lead to high variable outputs yield and all its components every year. These results are accepted by Moreno et al. (2003), which proceeds that the N fertilizer highly depends on growing season's variations conditioned by environmental factors. Climatic data of both growing season were indicated that the season of 2015/16 had favorable climate conditions for grain yield, without more cold in winter and good rainfall in planting time (October, November) for early germination and for grain filling time occurs (April, May). On the other hand, the climate conditions for all components without grain yield were good in 2016/17 growing season.

According to results of the study, the application of nitrogen doses had positive effect on yield and yield components. The results increased in parallel with dose increase for both cultivars for all characters. Nitrogen application had positive influence on all the yield components (Fallahi et al., 2008). The best grain yield and majority components results were obtained with N₃ nitrogen dosing. Moreover, N₄ and N₂ nitrogen dosing can be advice for some special components depend on climatic data of seasons. The results of nitrogen use efficiency of the present study agree with the findings of Janušauskaitė (2013) who reported that nitrogen use efficiency for grain yield and protein content as affected by nitrogen rate and its splitting in spring triticale and Yildirim et al. (2007) the findings suggest that advanced breeding lines should be selected at different N levels for better N use efficiency and genetic investigations

should be conducted in multiple environments and Alaru et al. (2009) who reported that yield values are under the influence of application of N fertilization. There are more N rate application studies on components in different crop as well as triticale which agree in present study; Hadi et al. (2012), increase in number of time to spike by increasing N rate and it might be attributed to the increase in long time filling grain (Gursoy, 2011; Shafi et al., 2011). Hadi et al. (2012), increase SS m⁻² by increasing N rate and it might be attributed to increase the time of number spike, Hadi et al. (2012), increase in PH by increasing N rate might be attributed to the increase in vegetative of plant. The increase SL depend on increasing N rate and available environmental in barley and wheat by researchers (Gursoy, 2011; Shafi et al., 2011). Subhan et al. (2004) and Shafi et al. (2011), increase NGS by increasing N rate and it might be attributed to increase the time grain occurs. Fallahi et al. (2008) Alazmani (2015), Yildirim et al. (2016), increasing N rate and it might be attributed to increase the grains per spike, and this is increasing GY.

GGE biplot analysis

The GGE biplot mainly allows the visualization of any crossover GE interaction, which is very important for the breeding program (Sayar, 2017; Kendal, 2015). The GGE biplot method has been widely used to analyze the stability and performance of the genotypes for yield and other components (Goyal et al., 2011; Tekdal and Kendal, 2017). Moreover, GEI and yield stability analyses are important for their consideration of both varietal stability and suitability for cultivation across seasons and ecological circumstances. The GT (genotype-trait) biplot provides an excellent tool for visualizing genotype × trait data (Kendal, 2015).

The AMMI analysis shows good visualization of interaction among nitrogen application doses and cultivars (*Fig. 1b*), among doses and years (*Fig. 1c*), among years and cultivars (*Fig. 1d*). The AMMI method provides considerable flexibility, allowing plant breeders to simultaneously select for yield and stability (Kendal and Sener, 2015).

The GGE biplot could be used to interpret the relationships among nitrogen doses, components, and groups of component (*Fig. 1e*). An understanding of the relationship between doses and components can aid in better understanding doses objectives and in identifying components that are positively or negatively correlated with nitrogen doses. This understanding can also aid in identifying components that can be indirectly selected by selecting for correlated components. It also helps to visualize the strengths and weaknesses of nitrogen doses, which is important for application in barley. If the angle of the vector was less than 90°, there was a positive correlation two observation factors. If the angle was equal to 90°, they were not correlated. There was a negative correlation if the angle was less than 90° (Yan and Thinker, 2006; Dogan et al., 2107). On the other hand, all components took place in two different groups, with N₂, N₃ and N₄ doses (*Fig. 1f*). There is high correlation which is took places in same group (Kendal et al., 2016). Moreover, the GGE biplot was accurate in interpreting the ranking and comparing N doses and components (*Fig. 1g*). The doses with both high mean performance and high stability for all of the components were called as table dose (N₂). The dose located side of some components, it means that this dose can be advise for these components (N₃ an N₄). The center of the concentric circles (i.e., ideal dose) was the AEA in the positive direction. In the comparison biplot doses located closer to the ideal dose were more desirable than others (Yan and Tinker, 2006; Benin et al., 2012; Dogan et al., 2016). The result demonstrated that N₂ was ideal application dose in the

both season for all components, as it was in the center circle for the ideal doses and on the AEA (Fig. 1h).

Conclusions

1. In the present study nitrogen application doses had positive effect on grain yield and yield components, and the values of component and grain yield increased in parallel with application nitrogen doses.

2. The nitrogen dose of N₃ (120 kg ha⁻¹) are responsible for the maximum productivity of triticale crop in Mardin province environmental conditions. Esin variety showed that it is best cultivar for all components and grain yield except PC.

3. On the other hand; AMMI and GGE biplot analysis revealed that this analysis provided useful results and high image quality to show the correlation among doses, cultivars and components.

4. The results of study showed that the N₃ application of doses in triticale should be recommend for next studies both farm and research area.

REFERENCES

- [1] Alaru, M., Laur, Ü., Eremeev, V., Reintam, E., Selge, A., Noormets, M. (2009): Winter triticale yield formation and quality affected by N rate, timing and splitting. – *Agricultural and Food Science* 18: 76-90.
- [2] Alazmani, A. (2015): Evaluation of yield and yield components of barley varieties to nitrogen. – *International Journal of Agriculture and Crop Sciences IJACS* 8(1): 52-54.
- [3] Anonymous (2002): SAS, Institute. JMP Statistics. – SAS Institute Inc., Cary, NC.
- [4] Asfaw, A., Alemayehu, F., Gurum, F., Atnaf, M. (2009): AMMI and SREG GGE biplot analysis for matching varieties onto soybean production environments in Ethiopia. – *Scientific Res. and Essay* 4(11): 1322-1330.
- [5] Benin, G., Bornhofen, E., Beche, E., Pagliosa, I. E. S., Lemesda, Silva, C., Pinnow, C. (2012): Agronomic performance of wheat cultivars in response to nitrogen fertilization levels. – *Acta Scientiarum, Agronomy Maringá* 34(3): 275-283.
- [6] Dogan, Y., Kendal, E., Oral, E. (2016): Identifying of relationship between traits and grain yield in spring Barley by GGE biplot analysis. – *Agriculture and Forestry* 6(24): 239-252.
- [7] Estrada-Campuzano, G., Slafer, G. A., Miralles, D. J. (2012): Differences in yield, biomass and their components between triticale and wheat grown under contrasting water and nitrogen environments. – *Field Crops Research* 128: 167-179. <http://dx.doi.org/10.1016/j.fcr.2012.01.003>.
- [8] Fallahi, H A, Nasserli, A., Siadat, A. (2008): Wheat yield components are positively influenced by nitrogen application under moisture deficit environments. – *Int. J. Agri. Biol.* 10: 6.
- [9] Farshadfar, E., Rashidi, M., Jowkar, M. M., Zali, H. (2013): GGE Biplot analysis of genotype × environment interaction in chickpea genotypes. – *European J. of Exper. Biology* 3(1): 417-423.
- [10] Goyal, A., Beres, B. L., Randhawa, H. S., Navabi, A., Salmon, D. F., Eudes, F. (2011): Yield stability analysis of broadly adaptive triticale germplasm in southern and central Alberta. – *Canadian J. Plant Sci.* 91: 125-135.
- [11] Gülmezoglu, N., Kinaci, E. (2004): Efficiency of different top dressed nitrogen on triticale (×*Triticosecale* Wittmack) under contrasting precipitation conditions in semiarid

- region. – Pakistan Journal of Biological Sciences 7(3): 353-358.
<http://dx.doi.org/10.3923/pjbs.2004.353.358>.
- [12] Gürsoy, M. (2011): Effects of different nitrogen doses on yield and quality components in some two-rowed barley (*Hordeum vulgare* L.) lines and cultivars. – Journal of New World Sciences 6(4): 114-123.
- [13] Hadi, F., Arif, M., Hussain, F. (2012): Response of dual purpose barley to rates and methods of nitrogen application. – ARPN Journal of Agricultural and Biological Science 7(7): 533-539.
- [14] Janušauskaitė, D. (2013): Spring triticale yield formation and nitrogen use efficiency as affected by nitrogen rate and its splitting. – Zemdirbyste-Agriculture 100(4): 383-392. DOI: 10.13080/z-a.2013.100.049.
- [15] Kendal, E. (2015): Relationship between chlorophyll and other features in durum wheat (*Triticum turgidum* L. var. durum) using SPAD and biplot analyses. – Journal of Agricultural Science and Technology 17: 1873-1886.
- [16] Kendal, E. (2016): GGE Biplot analysis of multi-environment yield trials in barley (*Hordeum vulgare* L) cultivars. – Ekin J. of Crop Breeding and Genetics 2(3): 90-99.
- [17] Kendal, E., Doğan, Y. (2015): Stability of a candidate and cultivars (*Hordeum vulgare* L) by GGE biplot analysis of multi-environment yield trial in spring barley. – Agriculture and Forestry/Poljoprivredai Sumarstvo 61: 4.
- [18] Kendal, E., Sayar, M. S. (2016): The stability of some spring triticale genotypes using biplot analysis. – The Journal of Animal and Plant Sciences 26(3): 754-765.
- [19] Kendal, E., Şener, O. (2015): Examination of genotype× environment interactions by GGE biplot analysis in spring durum wheat. – Indian J. Genet. 75(3): 341-348.
- [20] Kendal, E., Tekdal, S., Aktaş, H., Karaman, M. (2014): Determination compliance abilities of some triticale varieties and comparison with wheat in SoutheasternAnatolia conditions of Turkey. – Communications in Agricultural and Applied Biological Science 79(4): 192-200.
- [21] Kendal, E., Sayar, M. S., Tekdal, S., Aktaş, H., Karaman, M. (2016): Assessment of the impact of ecological factors on yield and quality parameters in triticale using GGE biplot and AMMI analysis. – Pak. J. Bot. 48(5): 1903-1913.
- [22] Kilic, H. (2014): Additive main effect and multiplicative interactions (AMMI) Analysis of grain yield in barley genotypes across environments. – J. Agr. Sc. 20: 337-344.
- [23] Lance, R. Gibson., Carasella, D., Nance, Douglas, L. K. (2007): Winter triticale response to nitrogen fertilization when grown after corn or soybean. – Agron. J. 99: 49-58.
- [24] Lestingi, A., Bovera, F., De Giorgio, D., Ventrella, D., Tateo, A. (2010): Effects of tillage and nitrogen fertilization on triticale grain yield, chemical composition and nutritive value. – Journal of the Science of Food and Agriculture 490: 2440-2446. <http://dx.doi.org/10.1002/jsfa.410>.
- [25] Moreno, A., Moreno, M. M., Ribas, F., Cabello, M. C. (2003): Influence of nitrogen fertilizer on grain yield of barley (*Hordeum vulgare* L.) under irrigated conditions. – Spanish Journal of Agricultural Research 1(1): 91-100.
- [26] Moselhy, E. I., Zahran, M. A. (2002): Effect of bio and mineral nitrogen fertilization on barley crop grown on a sandy soil. – Egypt. J. Agric. Res. 3: 921-936.
- [27] Sayar, M. S. (2017): Additive main effects and multiplicative interactions (AMMI) analysis for fresh forage yield in common vetch (*Vicia sativa* L.) genotypes. – Agriculture and Forestry 63(1): 119-127.
- [28] Sayar, M. S., Han, Y. (2015): Determination of seed yield and yield components of grasspea (*Lathyrus sativus* L.) lines and evaluations using GGE biplot analysis method. – TarımBilimleri Der- J. of Agric. Sci. 21(1): 78-92.
- [29] Shafi, M., Bakht, J., Jalal, F., Aman, Khan, M., Gül, Khattak, S. (2011): Effect of nitrogen application on yield and yield components of barley (*Hordeum vulgare* L.). – Pak. J. Bot. 43(3): 1471-1475.

- [30] Steel, G. D., Torrie, J. H. (1980): Principles and Procedures of Statistics: A Biometrical Approach. 2. ed. – McGraw-Hill, New York.
- [31] Subhan, W., Anwar, M., Ahmad, N., Gulzar, A., Siddig, M, Rahman, S., Ahmad, I, Rauf, A. (2004): Effect of gamma radiation on growth and yield of barley under different nitrogen levels. – Pakistan Journal of Biological Sci. 7(6): 981-983.
- [32] Ugarte, C., Calderini, D. F., Slafer, G. A. (2007): Grain weight and grain number responsiveness to pre-anthesis temperature in wheat, barley and triticale. – Field Crops Research 100: 240-248. <http://dx.doi.org/10.1016/j.fcr.2006.07.010>.
- [33] Yan, W., Tinker, N. A. (2006): Biplot analysis of multi-environment trial data. Principles and applications. – Canadian Journal of Plant Science 86: 623-645.
- [34] Yildirim, M., Yakut, Z., Akıncı, C., Kurt, F., Kizilgeci, F. (2016): Nitrogen rate and timing implementation on durum wheat in a bed planting system. – Sains Malaysiana 45(2): 221-228.
- [35] Yildirim, M., Bahar, B., Genc, I., Korkmaz, K., Karnez, E. (2007): Diallel analysis of wheat parents and their F2 progenies under medium and low level of available n in soil. – Journal of Plant Nutrition 30: 937-945.