

SEASONAL VARIATIONS IN SOIL CO₂ EMISSIONS UNDER CONTINUOUS FIELD CROP PRODUCTION IN SEMI ARID SOUTHEASTERN TURKEY

YILMAZ, G.

*Department of Environmental Engineering, Faculty of Engineering, Harran University, Osmanbey Campus, Şanlıurfa, Turkey
(e-mail: gyilmaz@harran.edu.tr; phone: +90-414-318-3789; fax: +90-414-318-3190)*

(Received 24th Jan 2019; accepted 29th Mar 2019)

Abstract. Quantification of soil CO₂ emission under different cropping designs and relating it to the climate parameters can help better understanding the main mechanisms controlling its magnitude and mitigating its release to the atmosphere. Wheat and corn are the most produced crops worldwide contributing highly to total CO₂ emissions from agricultural lands to atmosphere. Long term fluctuations in the soil CO₂ emissions under continuous wheat and corn cropping and bare land were monitored for 509 weeks from 2009 to 2018 in semi arid conditions in the Harran Plain located in southeastern Turkey. The soil CO₂ emissions were modeled for each cropping seasons separately using available meteorological parameters (air temperature (AT), soil temperatures (ST), relative humidity (RH) and rainfall) and field soil moisture (SM) data. The quantity of long term soil CO₂ emission were following order; corn, wheat and bare land and it was around 38% higher (574 vs 792 kg ha⁻¹ week⁻¹) in soils under field crops than bare soils. Overall there were n increasing trends in seasonal, yearly cumulative and average weekly soil CO₂ emissions over the years. The weekly soil CO₂ emission values were positively correlated with AT and ST but negatively with RH and SM. The linear models were able to explain variations in the long term weekly soil CO₂ emissions obtained during wheat, corn cultivations and bare land up to 77%, 55% and 50%, respectively.

Keywords: *global warming, soil CO₂ emission, wheat, corn, meteorological data, stepwise regression*

Introduction

Global warming increases the soil respiration causing the release of more CO₂ emissions (Lu and Cheng et al., 2009). Among other three greenhouse gases CO₂ has the highest concentration in the atmosphere. Measurements and modeling of the soil CO₂ emissions under different soil types and crop managements are cost effective and crucial for global warming, carbon cycling as well as international agreements that aim to mitigate the amount of carbon released to atmosphere (Allaire et al., 2012). Quantification of the CO₂ emission from soils under different land uses is also important for developing strategies for reducing the CO₂ emissions to atmosphere. A number of researchers have shown atmospheric CO₂ increase very recently. The soil CO₂ respiration consists of basically two parts; roots respiration and microbial respiration as a result of organic matter decomposition (Morell et al., 2012; Zhang et al., 2015). Roots and microbial activities are responsible of 40% and 60% of total the soil respiration, respectively (Kuzyokav and Larionova, 2006).

Methods such as edycovariance or with closed chamber combined with gas chromatography are generally used for measurement of the soil CO₂ emissions. Another most commonly used technique in measuring the soil CO₂ emissions is soda lime (Edwards, 1982). This is advantageous, because it is cheap and easy to perform, enabling the soil CO₂ measurements at higher number of locations simultaneously compare to aforementioned methods which can only be obtained only at limited number of locations.

In addition, the earlier researchers comparing soda lime method reported that there is no statistically significant different between soda lime method and them (Raich et al.1990).

There are nearly two times more carbon in the soils than atmosphere (Lehmann and Joseph, 2009) and according to Lal (2004) that may be more than that. Depending upon agricultural management, soils can either be source or sink for CO₂ in the atmosphere and the amount of CO₂ released from soils to atmosphere can change based on the soil type, land use type or vegetation type, agricultural management as well as the time of measurement (i.e. seasonal variability) (Iqbal et al., 2009; Panosso et al., 2009, 2011; Mancinelli et al., 2010). Soil CO₂ emissions in the agricultural soils are generally between 0.1 and 6 g CO₂ m⁻² h⁻¹ (Sauerbeck, 2001).

Average soil CO₂ emissions reported by earlier studies were 59 to 527 mg m² h⁻¹, 37-498 mg m² h⁻¹, 32-397 mg m² h⁻¹ for forest, grassland and cropland (Wang et al., 2008). The amount of soil CO₂ emissions in peat, sandy and clay soils respectively were obtained as 650 mg m² d⁻¹, 300 mg m² h⁻¹ and 500 mg m² h⁻¹ (Koizumi et al., 1999). Bauer et al. (2006) for the soils under conservation tillage measured soil CO₂ emissions as 840 mg m² h⁻¹. Akbolat et al. (2009) reported as 300-700 mg m² h⁻¹. The cropped soils release CO₂ 2 to 3 fold higher than bare soils (Rastogi, 2002).

The soil CO₂ emissions are mostly impacted from the factors such as soil temperature and moisture (Bauer et al., 2006; Ngao et al., 2012; Xu et al., 2016; Iqbal et al., 2009; Reth et al., 2005), weather parameters, soil organic matter, C/N ratio, bulk density (Lin et al., 2008; Morell et al., 2012).

Wheat and corn are the most produced grain crops in the world and Turkey is one of the largest producers of both. Yearly around 653 355 000 and 19 674 000 tonnes wheat is produced in 217 312 000 ha and 8 103 000 ha areas in the world and in Turkey, respectively (FAOSTAT, 2010). Worldwide wheat grows on 215 million hectares of land each year (CGIAR, 2017).

Emissions of CO₂ from most of the forest crops, orchards, grasses are well documented (Frank et al., 2006; Almaraz et al., 2009). On the other hand, soil CO₂ emissions from soils under field crops with long terms field studies are limited. The studies available are mostly performed in a short term period which may not be sufficient to construct reliable models. Therefore, information about soil CO₂ emissions of soils under wheat plantation makes it significant.

The goals of this study were to: 1) quantify weekly and seasonal distributions of the long terms (from 2009 to 2018) CO₂ emissions from the soil under wheat and corn plantation 2) compare CO₂ emissions estimations obtained from the soils under field conditions compared with bare soils located nearby field crops, and 3) model the relationships between the soil CO₂ emissions from the soils under different managements (wheat, corn cultivation and uncultivated bare soils) and meteorological parameters and field moisture data at the same locations with soil CO₂ measurements.

Materials and methods

Study site and field management

The study was performed in a field under continuous wheat (2009-2017) and corn (2016-2018) cultivations, in addition to bare land located in the experimental farm of the Harran University, southeastern Turkey (37° 10' N Lat; 38° 59' E Long; *Fig. 1*). The study area has a semi arid climate with an average annual temperature, precipitation, evaporation and altitude values of 17.2 °C, 365.2 mm, 1848 mm and 520 m, respectively.

The soils of the study area are high in percent clay content, pH, percent CaCO₃ content and cation exchange capacity (CEC) but low in soil organic matter (<1.5 g kg⁻¹) (Dinç et al., 1998). Detailed information about the soil properties of the area is given in *Table 1*. Soils were clay loam and classified as Vertisol, according to Soil Taxonomy (Dinç et al., 1998).

Table 1. Soil properties of bare field and the experimental areas under field crops

Location	Sampling depth (cm)	EC (dS/m)	CaCO ₃ (%)	pH	P	K	Organic matter	Total N (%)	Sand	Clay	Silt	Texture
Bare area	(0-30)	0.73	17.40	7.70	2.71	112.5	1.36	0.09	22	60	18	Clay
Bare area	(30-60)	0.82	17.40	7.76	0.82	82.0	0.85	0.08	22	60	18	Clay
Field crops	(0-30)	0.83	23.50	7.76	3.04	82.0	1.45	0.09	32	50	18	Clay
Field crops	(30-60)	0.82	25.40	7.84	0.41	36.70	1.01	0.09	26	56	18	Clay

Wheat crops were planted in 2009-2017. Also during the experiment, in 2016-2018, corn crop following wheat was planted. Wheat and corn received regular care which included application of water as flood irrigation, fertilizer and pest control with sowing, 40 kg da⁻¹ compose fertilizer as 20+20+Zn has been applied as base fertilizer and as top fertilizer 50 kg da⁻¹ urea was applied. The same fertilizer practice was repeated in each year.

Soil laboratory analysis

A composite soil sample was taken as representative of orchard field at the surface (0-30 cm) and subsurface soil (30-60, 60-90 cm). Soil samples were air dried and passed through a 2 mm sieve and analyzed according to the routine soil analyses methods. Particle size distributions were determined by the hydrometer method (Bouyoucos, 1926); organic matter content by the Walkley-Black method (Nelson and Sommers, 1982); CaCO₃ content using a calcimeter (Kacar, 1994); soil pH with a 1:2.5 from soil/water suspension using pH meter (McLean, 1982); electrical conductivity (EC) in soil extraction using a conductivity meter (Janzen, 1993), soil N according to Kjeldahl method (Kacar, 1994). Soil available P and K were determined using an ICP-AEAS (Varian-Vista, Palo Alto, CA, USA), after NaHCO₃ extraction (pH 8.5) (Olsen and Sommers, 1982) and NH₄OAc extraction (Knudsen et al., 1982), respectively.

Soil CO₂ emissions and meteorological variables

In addition to soil CO₂ emission measurements under wheat and corn plantation, the measurements were also performed in bare soils with no cultivation fields. The soil CO₂ emissions measurements were performed as weekly basis during total of 509 weeks between January, 2009 and October, 2018, using soda lime method (Edwards, 1982).

Soda lime method is advantageous because it is easy to perform and also does not require expensive equipments. Around 40 g soda lime was oven dried at 105 °C during 24 h, weighted and added in the numbered plastic PVC containers (3 cm tall, and 100 cm² area top). The containers that have a known amount of soda lime was placed under a chamber (23 cm width and 33 cm length) inserted 2 cm in soils in order to prevent interaction between soda lime and atmosphere in order to avoid any gas leakage underneath of the chamber as much as possible. A PVC lid (50 cm by 50 cm) was used to provide shadow for the chambers to reduce the temperature difference between inside

and outside of the chambers. After one week in the field, the chambers were left and PVC containers with soda lime were weighted in the laboratory to measure changes in weight during one week incubation period. The soil CO₂ emissions were performed with five replications and then averaged to obtain a final emission values.

The amount CO₂ emission was obtained as below:

$$F_{CO_2} = \frac{(C_{adsorbed} - C_{initial}) * W}{t_{incubation} * A} \quad (\text{Eq.1})$$

Here:

F_{CO₂} = emitted CO₂ g⁻¹ cm² h⁻¹

C_{adsorbed}: Weight of soda lime after incubation (g)

C_{initial}: Weight of soda lime before incubation (g)

W: coefficient of water, taken as 1.69

t_{incubation}: Incubation period (week)

A: Area of the chamber, cm²

Emission of CO₂ calculated as g cm² hour was converted into total weekly and monthly amounts.

Meteorological variables used in the study were obtained from Sanliurfa city meteorological station located around 5 km away from the study site. Daily measurements were converted to weekly basis. The meteorological variables included air temperature (AT), soil temperature (ST) at different depths (5, 10, 20, 50 and 100 cm), relative humidity (RH) and rainfall (mm). Field soil moisture data were obtained gravimetrically from the soils samples taken at the same location where soil CO₂ emissions were measured.



Figure 1. Photo showing the experimental design set up for soil CO₂ emissions

Statistical analyses and modeling

The relationships between CO₂ emissions at different sites (wheat and corn fields, and bare soils) and meteorological parameters and field soil moisture data were examined using correlation analyses and multiple linear regression (MLR) models.

MLR models the relationships between response (soil CO₂ emissions) and predictor variables (meteorological parameters and field soil moisture data) using a linear function between two (Eq. 2):

$$y_i = \beta_0 + \beta_1 X_{1i} + \dots + \beta_p X_{pi} + \varepsilon_i \quad (\text{Eq.2})$$

where y_i refers to response variable which is modeled by a linear function of predictor variables; X_i , β are coefficients; ε is error term.

The models were also validated using independent data set splitting the whole data into two groups; as 70 and 30%; while the former was used for calibration, the latter was used for validation of the model which was constructed using General Linear Model (GLM).

The models were constructed for the data in each sampling sites and each year separately and also for complete years (2009-2018). Correlation analyses and MLR modeling were performed using SPSS program. GLM model used in model validation (testing) was performed using R program.

Results

Soil properties

Averaged soil properties measured at different depths and sites are given in *Table 1*. Soils are high in CaCO₃ (>15%) and low in soil organic matter (<2%) contents. These were due to no organic residues given in to soils, conventional tillage and warm climate environment of the study area. Soils have a basic soil pH, a heavy clay texture and do not have any salinity problem which is indicated by low soil EC values (<4 dS m⁻¹). While total nitrogen and plant available P contents in the soils were low, K levels were high. Soil P and organic matter were higher in the soils under field crop rotation but it was opposite for K which was found higher in bare soils compare to the soils under wheat and corn crop rotation. It may be because of plant uptake.

Soil CO₂ emissions

Average, minimum and maximum of the soil CO₂ emissions (kg ha⁻¹ week⁻¹) measured in the soil during the study periods (2009–2018) are given in *Table 2*. Long term weekly, seasonal and cumulative yearly fluctuations of the soil CO₂ emissions are given in *Figures 2-5*. The results showed that CO₂ emissions were highly variable and the higher emission peaks were obtained in the summer seasons.

The soil CO₂ emissions were highly variable, both seasonally and locally ranging from 54 to 2504 kg ha⁻¹ week⁻¹. The CO₂ emissions in bare soils ranged from 57 to 1690 kg ha⁻¹ week⁻¹ with average values of 791 and 523 kg ha⁻¹ week⁻¹, respectively (*Table 2*). When the fields are under either wheat or corn plantation (excluding the weeks when there is no crop in the field), average soil CO₂ emissions were 534 and 850 kg ha⁻¹ week⁻¹ for bare soils (CO₂-Bare) and field soils (CO₂-Field), respectively.

Average weekly CO₂ emissions of the soils under wheat plantations in 2009-2017 were 310, 480, 516, 481, 400, 535, 680 and 696 kg ha⁻¹ week⁻¹, respectively, which showed a consistent increase over time. Similar to the soils under wheat plantations, average weekly CO₂ emissions of soils under corn plantation were higher compared to the previous years. It was 914 and 1056 kg ha⁻¹ week⁻¹ in 2016 and 2017, respectively. The average weekly soil CO₂ emissions from bare soils (CO₂-Bare) were always lower than those obtained in the soils under both wheat and corn cultivation (Table 2). Average weekly soil CO₂ emissions from the bare soils corresponding to the same weeks when fields under wheat plantation were 310, 480, 516, 481, 400, 535, 680 and 696 kg ha⁻¹ week⁻¹, respectively in 2009-2017. Average weekly soil CO₂ emissions from bare soils corresponding to the same weeks when the fields under corn plantation in 2016 and 2017 were 687 and 621 kg ha⁻¹ week⁻¹, respectively.

Table 2. Descriptive statistics of long term soil CO₂ emissions under different field management and meteorological parameters

		CO ₂ -B	CO ₂ -F	SM-B	SM-F	AT	RH	ST5	ST10	ST20	ST50	ST100	Rainfall
		kg ha ⁻¹ week ⁻¹		%		°C	%	°C					mm
2009 WHEAT (n†=25)	Avg.	310	506	24	29	15	53	16	16	15	15	15	8.1
	Min.	93	195	9	10	2	21	3	4	5	8	11	0.0
	Max.	505	1041	33	49	30	80	33	31	29	28	25	38.6
2010 WHEAT (n=34)	Avg.	480	575	28	24	15	56	16	16	16	16	17	10.7
	Min.	257	274	10	5	5	25	6	6	7	10	12	0.0
	Max.	860	1000	36	35	30	85	35	33	30	28	26	57.0
2011 WHEAT (n=28)	Avg.	516	1128	27	24	14	55	15	15	15	15	15	12.8
	Min.	366	275	9	6	5	31	6	7	8	9	11	0.0
	Max.	1015	2316	34	36	29	79	29	29	27	27	23	57.2
2012 WHEAT (n=29)	Avg.	481	786	27	31	14	50	14	14	14	14	14	14.3
	Min.	68	54	14	21	3	15	5	5	6	7	9	0.0
	Max.	831	1476	42	42	33	87	34	32	30	29	24	84.7
2013 WHEAT (n=30)	Avg.	400	853	28	30	14	58	14	14	14	15	15	14.6
	Min.	225	278	13	20	3	24	6	6	7	8	11	0.0
	Max.	746	1396	37	37	28	85	28	27	26	26	22	52.0
2014 WHEAT (n=23)	Avg.	535	769	24	22	16	47	16	16	15	15	15	8.2
	Min.	223	421	8	6	7	28	7	7	8	8	11	0.0
	Max.	783	1144	39	36	26	82	29	29	27	27	23	74.2
2016 WHEAT (n=23)	Avg.	680	886	23	24	15	51	16	16	16	15	15	7.3
	Min.	364	335	6	7	1	26	5	6	7	7	10	0.0
	Max.	992	1941	40	40	28	79	29	29	28	27	24	36.6
2016 CORN (n=22)	Avg.	687	914	12	14	27	33	29	29	28	28	27	1.8
	Min.	334	243	3	4	12	19	12	12	14	15	19	0.0
	Max.	1690	2504	33	38	34	66	36	35	34	34	30	39.6
2017 WHEAT (n=24)	Avg.	696	952	22	26	15	48	16	16	16	16	15	6.2
	Min.	512	550	10	7	4	25	6	6	6	7	9	0.0
	Max.	945	1327	31	37	30	73	32	31	30	29	25	60.8
2017 CORN (n=25)	Avg.	621	1056	12	31	21	43	23	23	23	23	24	2.9
	Min.	396	548	5	4	6	22	7	8	10	10	14	0.0
	Max.	911	1854	24	51	34	72	36	36	34	33	30	27.9
PLANTED & UNPLANTED (n=509)	Avg.	524	792	19	21	20	45	21	21	20	20	20	7.1
	Min.	57	54	1	1	1	15	3	4	5	7	9	0.0
	Max.	1690	2504	44	51	36	88	41	39	36	34	32	89.4
PLANTED (n=289)	Avg.	534	850	23	26	17	49	19	18	18	18	18	8.5
	Min.	68	54	3	4	1	15	3	4	5	7	9	0.0
	Max.	1690	2504	42	51	34	87	38	36	34	34	31	84.7

†The number of total weeks between planting and harvesting of with corresponding field crop




Figure 2. The weekly fluctuations of field soil CO₂, bare soil CO₂ emissions and air temperature from 2009 to 2018 (509 weeks)




Figure 3. Changes in yearly mean soil CO₂ emissions from two different sites and air temperature




Figure 4. Variations in cumulative yearly soil CO₂ emissions




Figure 5. Seasonal (each points corresponding to seasons starting from spring and followed by summer-fall-winter seasons) variations in soil CO₂ emissions from 2009 to 2018

Correlations between the soil CO₂ emissions from the bare soils and wheat and corn plantations for each year separately and combined are given in *Table 3*. When all years were considered together, weekly CO₂-field values were significantly correlated ($p < 0.05$) with all meteorological parameters except rainfall, soil temperature at 100 cm and soil moisture data. While weekly CO₂-bare soil values were significantly correlated ($p < 0.05$) with all meteorological and soil moisture parameters (*Table 3*).

In the 2012, 2013, 2016 and 2017 years, weekly soil CO₂ emission values obtained had a statistically significant positive correlations ($p < 0.05$) with air and soil temperatures at different depths while significant negative correlations ($p < 0.05$) with relative humidity. They had statistically significant negative correlation with soil moisture values in 2012 and 2013 and insignificant negative correlations in 2016 and 2017.

In the 2009 and 2011 years, only weekly soil CO₂ emission values obtained from the soils under wheat cultivation had statistically significant negative and positive correlations ($p < 0.05$) with the same variables. The same situation was valid for only weekly soil CO₂ emission values obtained from the bare soils in 2014 (*Table 3*). Rainfall values had statistically ($p < 0.05$) significant negative correlations with weekly soil CO₂ emission values obtained from the soils under wheat cultivation in 2010, 2012 and 2016 years (*Table 3*).

Correlations between the soil CO₂ emissions under corn plantation both in 2016 and 2018 and environmental parameters were mostly insignificant ($p > 0.05$) (*Table 3*). As opposed to bare land and soils under wheat cultivation, the soil CO₂ emissions measured during corn plantation in 2016 had negative correlations with air and soil temperatures but positive correlations with rainfall, relative humidity and soil moisture.

Modeling of soil CO₂ emissions

Multiple Linear Regression models were constructed between the soil CO₂ emissions at different locations (bare land and field crops) and available climate parameters; air

temperature, soil temperature, relative humidity, precipitation and field soil moisture data. For different land managements the models were constructed for all weeks in whole years (from 2009 to 2018), for only weeks when the field is planted with wheat or corn (from 2009 and 2018) and for weeks during crop season in each year, separately. Coefficient of Determination (R²) values showing the per cent variations in soil CO₂ emissions at different time and locations explained by each models and significance of independent variables in terms of contribution to the model are given in Table 4. Model R² values for complete years considering both planted and also unplanted weeks and considering only planted weeks were 0.16 and 0.22, respectively. They were 0.1 and 0.23 respectively for bare land. For the models constructed separately for each crop season in different years, R² values ranged from 0.47 to 0.77 for wheat plantation, between 0.19 and 0.55 for corn plantation and 0.12 to 0.66 for bare land (Table 4).

Table 3. Correlations between soil CO₂ emissions and climatic variables and soil moisture field data

		CO ₂ -F	SM-B	SM-F	AT	RH	ST5	ST10	ST20	ST50	ST100	Rainfall
		kg ha ⁻¹ week ⁻¹	%		°C	%	°C					mm
2009 WHEAT	CO2-B	0.48*	-0.36	-0.20	0.40*	-0.32	0.36	0.36	0.35	0.35	0.36	0.21
	CO2-F	-	-0.37	-0.29	0.59**	-0.55**	0.59**	0.58**	0.58**	0.58**	0.56**	-0.09
2010 WHEAT	CO2-B	0.82**	-0.18	-0.24	0.14	-0.33	0.18	0.19	0.18	0.16	0.13	-0.28
	CO2-F	-	-0.17	-0.28	0.22	-0.34*	0.26	0.27	0.26	0.22	0.16	-0.37*
2011 WHEAT	CO2-B	0.13	-0.10	-0.07	0.28	-0.17	0.26	0.27	0.28	0.30	0.42*	0.06
	CO2-F	-	-0.01	0.20	0.81**	-0.52**	0.83**	0.82**	0.81**	0.80**	0.67**	-0.13
2012 WHEAT	CO2-B	0.75**	-0.43*	-0.38*	0.59**	-0.66**	0.55**	0.54**	0.53**	0.52**	0.46*	-0.37*
	CO2-F	-	-0.46*	-0.47**	0.61**	-0.56**	0.59**	0.59**	0.58**	0.56**	0.47**	-0.40*
2013 WHEAT	CO2-B	0.49**	-0.75**	-0.69**	0.46*	-0.39**	0.52**	0.53**	0.52**	0.52**	0.44*	-0.26
	CO2-F	-	-0.61**	-0.63**	0.76**	-0.59**	0.75**	0.76**	0.75**	0.74**	0.63**	-0.33
2014 WHEAT	CO2-B	0.34	-0.22	-0.38	0.55**	-0.37	0.51*	0.52*	0.52*	0.52*	0.50*	-0.13
	CO2-F	-	0.40	0.52*	-0.36	0.36	-0.40	-0.40	-0.39	-0.39	-0.36	-0.15
2016 WHEAT	CO2-B	0.18	-0.31	-0.42*	0.51*	-0.58**	0.53**	0.53**	0.52*	0.52*	0.48*	-0.24
	CO2-F	-	-0.34	-0.27	0.55**	-0.47*	0.55**	0.54**	0.54**	0.53**	0.48**	-0.45*
2017 WHEAT	CO2-B	0.69**	-0.38	-0.35	0.44*	-0.57**	0.41*	0.41*	0.41*	0.40	0.38	-0.22
	CO2-F	-	-0.35	-0.32	0.57**	-0.29	0.56**	0.56**	0.56**	0.55**	0.53**	0.16
2016 CORN	CO2-B	0.70**	-0.03	0.02	-0.23	-0.04	-0.26	-0.26	-0.27	-0.28	-0.25	-0.07
	CO2-F	-	-0.11	0.05	-0.34	0.10	-0.32	-0.31	-0.31	-0.31	-0.25	0.01
2017 CORN	CO2-B	0.36	-0.22	-0.51**	0.18	-0.21	0.19	0.20	0.20	0.21	0.24	-0.09
	CO2-F	-	-0.63**	-0.25	0.58**	-0.48*	0.61**	0.62**	0.63**	0.64**	0.68**	-0.15
2018 CORN	CO2-B	0.36	0.02	0.07	-0.20	-0.01	-0.27	-0.28	-0.29	-0.30	-0.32	0.12
	CO2-F	-	-0.58**	0.40*	0.29	-0.23	0.29	0.29	0.30	0.31	0.34	-0.12
PLANTED & UNPLANTED	CO2-B	0.51**	-0.15**	-0.10*	0.17**	-0.21**	0.15**	0.15**	0.16**	0.15**	0.13**	-0.14**
	CO2-F	-	0.01	0.14**	0.14**	-0.10*	0.12**	0.12**	0.12**	0.11*	0.06	-0.07
PLANTED	CO2-B	0.49**	-0.37**	-0.31**	0.31**	-0.40**	0.30**	0.30**	0.31**	0.31**	0.30**	-0.24**
	CO2-F	-	-0.30**	-0.07	0.40**	-0.36**	0.38**	0.38**	0.38**	0.38**	0.35**	-0.19**

AT: Air temperature (°C); ST5, ST10 and ST100 are soil temperature at 5, 10 and 100 cm, respectively; SM: Soil Moisture; RH: Relative; * significant at p < 0.05 level; ** significant at p < 0.01 level

Both SM and ST at different depths were found significant (p < 0.05) in the models constructed for the soils under field crops (Table 4). For models constructed for soil CO₂ emissions in bare land, the ST20 and ST100 were significant when considering

whole weeks (planted and unplanted), the SM, RH and ST5 were statistically significant when considering only weeks in crop seasons (planted) (Table 4).

Table 4. MLR modeling results for soil CO₂ emissions under different land managements

	2009 Wheat	2010 Wheat	2011 Wheat	2012 Wheat	2013 Wheat	2014 Wheat	2016 Wheat	2017 Wheat	2016 Corn	2017 Corn	2018 Corn	Planted & Unplanted	Planted
CO ₂ -FIELD													
R2	0.54	0.47	0.77	0.53	0.60	0.66	0.60	0.48	0.19	0.55	0.27	0.16	0.22
SM-F	ns	ns	ns	ns	ns	**	ns	ns	ns	ns	ns	**	*
AT	ns	**	ns	ns	ns	**	*	ns	ns	ns	ns	ns	ns
RH	ns	ns	ns	ns	ns	*	*	ns	ns	ns	ns	ns	ns
ST5	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	*
ST10	ns	ns	ns	ns	ns	s	ns	ns	ns	ns	ns	ns	ns
ST20	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	**	*
ST50	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
ST100	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	*
Rainfall	ns	ns	ns	ns	ns	*	ns	ns	ns	ns	ns	ns	ns
CO ₂ -BARE													
R2	0.55	0.48	0.43	0.54	0.62	0.50	0.66	0.47	0.28	0.12	0.34	0.07	0.23
SM-B	ns	ns	ns	ns	**	ns	ns	ns	ns	ns	*	ns	**
AT	ns	**	ns	ns	ns	*	ns	ns	ns	ns	ns	ns	ns
RH	ns	*	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	**
ST5	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	**
ST10	ns	*	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
ST20	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	**	ns
ST50	ns	ns	ns	ns	ns	ns	*	ns	ns	ns	ns	ns	ns
ST100	ns	ns	*	ns	ns	ns	*	ns	ns	ns	ns	**	ns
Rainfall	ns	ns	ns	ns	ns	ns	*	ns	ns	ns	ns	ns	ns

R²: coefficient of determination; AT: air temperature (°C); ST5, ST10 and ST100 are soil temperature at 5, 10 and 100 cm, respectively; SM: soil moisture; RH: relative humidity

For the models constructed for each crop seasons in different years separately, the significance of independent variables in the models changed based on different crop types and also years. Air temperature were found significant only in the models constructed for soils under wheat plantation in 2010, 2014 and 2016 years and for bare soils in 2010 and 2014 years. Relative humidity values were significant only for models constructed for soils under wheat plantation in 2014 and 2016 and bare soils in 2019. ST at different depths were not found significant in the models constructed for each cropping season year separately for field crops but found significant for bare soils in 2010 (ST10), 2011 (ST100) and 2016 (ST50 and ST100) years. Rainfall was found statistically significant for the models constructed for soils under wheat plantation in 2014 and for bare land in 2016 cropping season (Table 4).

Stepwise regression was used to select the most important variables in modeling the soil CO₂ emissions in different cropping seasons and managements. The most significant variables selected by stepwise regressions are shown in Table 5. Accordingly, when all weeks (planted and unplanted weeks) considered, the method selected SM and AT as the most important variables explaining the variations in the soil CO₂ emissions under field crops and RH in soil CO₂ emissions from bare land. When considering only planted weeks, the result did not change for soils under field crops but

SM, ST5 and ST10 variables in addition to RH were also found as the most significant in modeling the soil CO₂ emissions from bare land.

Table 5. Significant meteorological parameters in modeling soil CO₂ emissions determined by stepwise regression

		VARIABLES	R ²	P
2009 WHEAT	CO ₂ -FIELD	AT	0.35	**
	CO ₂ -BARE	AT	0.16	*
2010 WHEAT	CO ₂ -FIELD	Rainfall	0.14	*
	CO ₂ -BARE	-	-	-
2011 WHEAT	CO ₂ -FIELD	ST5, SM-Field	0.75	**
	CO ₂ -BARE	ST100	0.18	*
2012 WHEAT	CO ₂ -FIELD	AT	0.38	**
	CO ₂ -BARE	RH	0.44	**
2013 WHEAT	CO ₂ -FIELD	AT	0.57	**
	CO ₂ -BARE	SM-Bare	0.57	**
2014 WHEAT	CO ₂ -FIELD	SM-Field	0.27	*
	CO ₂ -BARE	AT	0.31	**
2016 WHEAT	CO ₂ -FIELD	AT	0.31	**
	CO ₂ -BARE	RH	0.33	**
2017 WHEAT	CO ₂ -FIELD	AT	0.32	**
	CO ₂ -BARE	RH	0.32	**
2016 CORN	CO ₂ -FIELD	-	-	-
	CO ₂ -BARE	-	-	-
2017 CORN	CO ₂ -FIELD	ST100	0.46	**
	CO ₂ -BARE	-	-	-
2018 CORN	CO ₂ -FIELD	SM-Field	0.16	*
	CO ₂ -BARE	-	-	-
PLANTED	CO ₂ -FIELD	AT,SM-Field,	0.18	**
	CO ₂ -BARE	RH,SM-Bare,ST5,ST10	0.21	**
PLANTED & UNPLANTED	CO ₂ -FIELD	SM-Field, AT,	0.10	**
	CO ₂ -BARE	RH	0.04	**

R²: coefficient of determination; *p significant at p < 0.05 level; **significant p < 0.01 level
 Variables selected according to stepwise regression; AT: air temperature (°C); ST5, ST10 and ST100 are soil temperatures at 5, 10 and 100 cm, respectively; SM: soil moisture; RH: relative humidity

In other stepwise regression models performed for each crop season separately, the model selected AT as the most significant variable in 2009, 2012, 2013, 2016 and 2017 in explaining the variations in CO₂ emissions from the soils under wheat cultivation, in 2009 and 2014 years from bare land. Rainfall was only found as the most significant parameter in explaining the variations in CO₂ emissions from the soils under wheat cultivation in 2010, when stepwise regression is used. The RH was found the most significant parameter for the variations in soil CO₂ emissions from bare land. Soil temperatures at 5 cm and 100 cm were found the most significant variables for the variations in CO₂ emissions in 2011 from the soils under wheat cultivation and bare land. The ST100 was also found the most significant variable for the variations in soil

CO₂ emissions from the corn field in 2017. Soil moisture was selected as the most significant variable by stepwise regression in explaining variations in the soil CO₂ emissions from wheat plantation in 2011 and 2014 years and from bare land in 2013 (Table 5).

Seasonal variations in the soil CO₂ emissions from both field and also bare land were also estimated using GLM model and available meteorological parameters and soil moisture data. For this purpose, the data was first split into two groups. 70% of the data was used for model construction and the rest 30% was estimated using constructed model. R² and RMSE values showing the accuracy of the estimations of soil CO₂ emissions of the samples in validation set (test set) were 0.15 and 0.05 and 253 and 178 kg ha⁻¹ week⁻¹ for field and bare area, respectively. Figure 6 shows the consistency between actual and model predictions for the CO₂ values in both locations.




Figure 6. Actual and GLM model predictions of soil CO₂ emissions in field (CO₂-FIELD; R²: 0.148; RMSE: 253) and bare areas (CO₂-BARE; R²: 0.05; RMSE: 178)

Discussion

Around 20% of atmospherically CO₂ is due to agricultural activities (Lal and Bruce, 1999) and grain crops such as wheat and corn are the most produced grain crops worldwide therefore the quantification of soil CO₂ emissions under these types of vegetation are important in terms of accuracy of carbon budgets and modeling.

Long term weekly soil CO₂ emission amounts from field crops and bare land had a distinct trend in seasonal fluctuation. Higher in summer and lower in winter seasons similar to air temperature (*Fig. 2*). Seasonal trends have been often reported for CO₂ emissions from soil under various other crops (Raich et al., 2002).

Soil CO₂ emissions under field crops showed a clearly higher release compare to bare soils. The relation between soil CO₂ emissions over the years and mean air temperature can be observed in *Figure 3*.

The soil CO₂ emissions showed an increase over time, as can be observed in *Figures 4* and *5*. The yearly cumulative soil CO₂ emissions could be explained with increased temperature over years. In addition, the plant residuals increased causes increase in soil organic matter. In some years the rates showed a decrease compared to previous year, however there was a general trend toward the higher cumulative emission.

Soil CO₂ emissions measured varied highly based on time of measurement (seasonal-climate), management (field crop vs. bare land) and cropping season (type of crop). Guo et al. (2013) also showed that the quantity of the soil CO₂ emissions under field cropping (wheat) can change depending on different growing stage of the crop in addition to environmental parameters such as soil temperature and rainfall.

The soil CO₂ emissions obtained from field crop plantation of our study were in a similar range with other ecosystems. Meng-yang et al. (2014) reported soil CO₂ emissions as 829, 629 and 474 g CO₂ m⁻² from July 1 to September 30 (921, 698 and 526 kg ha⁻¹ week⁻¹, respectively) for fields under 20- year continuous maize, wheat and soybean cultivation in a similar semi arid area located in northeast China. Gu et al. (2016) reported the soil CO₂ emissions under wheat plantation ranging from 372 to 418 mg m⁻² h⁻¹ (equals to 625-702 kg ha⁻¹ week⁻¹). Chen et al. (2004) indicated soil CO₂ emissions in wheat field ranging from 10 to 606 mg m⁻² h⁻¹ (equals to 17-1019 kg ha⁻¹ week⁻¹). The average soil CO₂ emissions from under semi-arid continuous wheat was 2.34 μ mol m⁻² s⁻¹ (equals to 622 kg ha⁻¹ week⁻¹) by Zhang et al. (2011).

The CO₂ emissions obtained in our study were relatively lower than other ecosystems such as orchards, forests or pastures. In arid region in China for grassland in the Black Chinese soils, Li et al. (2009) measured average weekly soil CO₂ emissions during growing season as 1680 kg ha⁻¹ week⁻¹. Lessard et al. (1994) measured the soil CO₂ emissions under forest vegetation and found a range between 158 and 1037 kg ha⁻¹ week⁻¹ and also compared with nearby cultivated fields, which provided soil CO₂ emissions between 18 and 495 kg ha⁻¹ week⁻¹. This is three times lower than emissions in the soils under forest. As opposed to these researchers, Iqbal et al. (2009) comparing the soil CO₂ fluxes under different land management, such as vegetables, forest, uplands and orchard stated that the soils under forest and orchard management provided relatively lower CO₂ emissions compare to other agricultural land uses.

According to average weekly emissions, the soils in continuous wheat and corn emitted 38% more CO₂ than the bare soils. The CO₂ emission from soil is the combination of root and soil microbial activities (Kuzyakov and Larionova, 2006). The root contributions to soil emissions are known as 48% (Raich and Tufekcioglu, 2000).

Soils under plantation can provide 2 to 3 fold higher CO₂ emissions compared to bare soil.

When considering the years for both corn and wheat; the mean soil CO₂ emissions measured during corn production was higher than wheat production in both 2016 and 2017. This may be due to differences in time of growing wheat and corn. Wheat grows in winter and spring seasons, corn is grown in summer season, mostly when the air and soil temperatures are higher. The researcher who compared the soil CO₂ emissions from wheat and corn cultivation in a field experiment specifically designed for comparison purpose reported that emissions were higher in corn than both wheat and also soybean cultivations. The researcher attributed the increase to that of higher residues incorporated into the soil in corn cultivation. Our study was completely under farmer conditions with regular field cares. Relatively higher emissions from the soil under corn crops have been attributed to higher residue incorporation under corn vegetation compare to other crops compared such as wheat and soybean.

We found statistically significant correlations between the soil CO₂ emissions and meteorological variables available and field soil moisture data. Except a few cases, the soil CO₂ emissions had mostly significant positive correlations with AT and ST while significant negative correlations found with RH, rainfall and soil moisture. Significance of variables such as the soil temperature and moisture in modeling the soil CO₂ emissions have been mostly emphasized by earlier researchers. The negative correlations between soil the CO₂ emissions and soil moisture content and rainfall could be due to optimum moisture conditions during the study period which was provided by irrigation. Earlier researchers reported that as long as soil moisture are at optimum level, its impact may be insignificant in the soil CO₂ emissions (Morell et al., 2010; Mancinelli et al., 2010).

Modeling of soil CO₂ emissions

Although the models used to estimate the soil CO₂ emissions at different locations, crop managements and seasons were found statistically significant ($p < 0.05$). The R² values showing capability of MLR models for explaining variations in the long terms soil CO₂ emissions were relatively lower, indicating that there are other factors that may involve in overall variations in the emissions. Zhang et al. (2011) emphasized that soil management practices such as tillage had high control over the quantity of soil CO₂ emissions under field conditions in addition to environmental factors such as soil temperature and moisture.

Lower R² values have been also found by earlier researchers. Mapanda et al. (2010) obtained R² values between the soil CO₂ emissions and various environmental parameters were between 0.12 and 0.48 from the models. Mancinelli et al. (2010) received R² values between 0.25 and 0.28 using the soil temperature and moisture as controlling parameter.

According to the MLR and stepwise regression models, the significant and most significant parameters selected in modeling relationships between long terms fluctuations in soil CO₂ emissions from different crop management and climatic parameters along with field soil moisture data were not same for the models constructed for different seasons and cropping managements. All variables included into the models were once selected as the most significant variables in modeling, however in most cases, AT was selected as the most significant variable in modeling the soil CO₂

emissions, which was followed by soil moisture, soil temperatures at different depth, RH and rainfall data.

Overall the accuracies of the soil CO₂ estimations using GLM models and available meteorological data were low. However the model estimations closely followed the actual observations (*Fig. 6*). Low estimation accuracies were mostly due to the higher emission values within the data sets which were not being able to be captured by the model and were underestimated. In addition to climate variables, agricultural practices such as plowing which are not included in the model may impact magnitude of the distribution of soil CO₂ emissions and cause occurrence of extreme values.

Conclusions

This study evaluated the long term (from 2009 to 2018) seasonal fluctuations in the soil CO₂ emissions under wheat and corn plantation and compared their results with ones obtained from bare soils located nearby. The CO₂ emissions from the soils under field crops were always about 38% higher than bare soils. The soil CO₂ emissions showed an increasing trend over the years which can be explained by increases in air temperature and accumulations of plant residues in the soil. Up to 77% of variations in the soil CO₂ emissions were explained with combined use of MLR model and available meteorological parameters and soil moisture data. Air temperature, soil temperatures at different depth and soil moisture were found the most significant parameters impacting the overall variations in long terms fluctuations in the soil CO₂ emissions under different cropping and bare soils. Overall the results indicated that per cent of variations explained by modeling could be improved by incorporating the soil management factors into models.

Acknowledgements. Financial support of this study has been provided by Harran University Scientific Research Commissions (HUBAK). The author would like to thank Prof. Dr. Ahmet Ruhi Mermut for his suggestion to improve the text and Dr. A. V. Bilgili, in Harran University, Turkey, for his contributions to the manuscript preparation and support.

REFERENCES

- [1] Akbolat, D. Evrendilek, F., Coskan, A., Ekinçi, K. (2009): Quantifying soil respiration in response to shortterm tillage practices: a case study in southern Turkey. – *Acta Agriculturae Scandinavica Section B-Soil and Plant Science* 59: 50-56.
- [2] Almaraz, J. J., Mabood, F., Zhou, X., Madramootoo, C., Rochette, P., Ma, B., Smith, D. L. (2009): Carbon dioxide and nitrous oxide fluxes in corn grown under two tillage systems in southwestern Quebec. – *Soil Science Society of America Journal* 73: 113-119.
- [3] Allaire, S. E., Lange, S. F., Lafond, J. A., Pelletier, B., Cambouris, A. N., Dutilleul, P. (2012): Multiscale spatial variability of CO₂ emissions and correlations with physico-chemical soil properties. – *Geoderma* 170: 251-260.
- [4] Bauer, P. J., Frederick, J. R., Novak, J. M., & Hunt, P. G. (2006): Soil CO₂ flux from a Norfolk loamy sand after 25 years of conventional and conservation tillage. – *Soil & Tillage Research* 90: 205-211.
- [5] Bouyoucos, G. J. (1926): Estimation of the colloidal material in soils. – *Science* 64: 362.
- [6] CGIAR (2017): 2016 Annual Report to CGIAR Consortium. Copenhagen, Denmark: CGIAR Research Program on Climate Change. – Agriculture and Food Security (CCAFS), Wageningen.

- [7] Dinç, U., Şenol, S., Sayın, M., Kapur, S., Güzel, N., Dericci, R., Yeşilsoy, M. Ş. et al. (1988): Güneydoğu Anadolu Bölgesi Toprakları (GAP), I. Harran Ovası (in Turkish). – TÜBİTAK, Şanlıurfa (Project No: TOAG-534).
- [8] Edwards, N. T. (1982): The use of soda-lime for measuring respiration rates in terrestrial systems. – *Pedobiologia* 23: 321 – 330.
- [9] FAOSTAT (2010): Food and Agricultural Organization of the United Nations, Rome. – FAOSTAT@fao.org.
- [10] Frank, A. B., Liebig, M. A., Tanaka, D. L. (2006): Management effects on soil CO₂ efflux in northern semiarid grassland and cropland. – *Soil & Tillage Research* 89: 78-85.
- [11] Guo, Q., Li, W. W., Liu, D. D., Wu, W., Liu, Y., Wen, X. X., Liao, Y. C. (2013): Seasonal characteristics of CO₂ fluxes in a rain-fed wheat field ecosystem at the Loess Plateau. – *Spanish Journal of Agricultural Research* 11: 980-988.
- [12] Iqbal, J., Lin, S., Hu, R., Feng, M. (2009): Temporal variability of soil-atmospheric CO₂ and CH₄ fluxes from different land uses in mid-subtropical China. – *Atmospheric Environment* 43: 5865-5875.
- [13] Janzen, H. H. (1993): Soluble Salts. – In: Carter, M. R. (ed.) *Soil Sampling and Methods of Analysis*. – Canadian Society of Soil Science, Crc Pres Inc. Boca Raton, FL.
- [14] Kacar, B. (1994): *Soil and Plant Analysis III. Soil Analysis*. – Ankara Uni. Agricultural Fac., Ankara.
- [15] Knudsen, D., Peterson, G. A., Pratt, P. F. (1982): Lithium, Sodium, and Potassium. – In: Page, A. L., Miller, R. H., Keeney, D. R. (eds.) *Methods of Soil Analysis, Part 2*. ASA, Madison, WI.
- [16] Koizumi, H., Kontturi, M., Mariko, S., Nakadai, T., Bekku, Y., Mela, T. (1999): Soil respiration in three soil types in agricultural ecosystems in Finland. – *Acta Agriculturae Scandinavica, Section B—Soil & Plant Science* 49: 65-74.
- [17] Kuzyakov, Y. V., Larionova, A. A. (2006): Contribution of rhizomicrobial and root respiration to the CO₂ emission from soil (A review). – *Eurasian Soil Sci.* 39: 753.
- [18] Lal, R., Bruce, J. P. (1999): The potential of world cropland soils to sequester C and mitigate the greenhouse effect. – *Environmental Science & Policy* 2: 177-185.
- [19] Lehmann, J., Joseph, S. (2009): *Biochar for Environmental Management: Science and Technology*. – Earthscan, London.
- [20] Lessard, R., Rochette, P., Topp, E., Pattey, E., Desjardins, R. L., Beumont, G. (1994): Methane and carbon dioxide fluxes from poorly drained adjacent cultivated forest sites. – *Can J Soil Sci* 74: 139-146.
- [21] Li, H., Han, X., Qiao, Y., Ho, X., Xing, B. (2009): Carbon dioxide emission from black soil as influenced by land-use change and long-term fertilization. – *Commun in Soil Sci Plan Analysis* 40: 1350-1368.
- [22] Lin, T. B., Wang, Z. Q., Song, X. L., Qu, Y. W., Meng, Z. Y. (2008): CO₂ flux and impact factors in winter wheat field ecosystem. – *Chin Eco Agr* 16: 458-1463.
- [23] Lu, X., Cheng, G. (2009): Climate change effects on soil carbon dynamics and greenhouse gas emissions in *Abies fabri* forest of subalpine, southwest China. – *Soil Biology and Biochemistry* 4: 1015-1021.
- [24] Lu, X., Lu, X., Tanveer, S. K., Wen, X., Liao, Y. (2016): Effects of tillage management on soil CO₂ emission and wheat yield under rain-fed conditions. – *Soil Research* 54: 38-48.
- [25] Mancinelli, R., Campiglia, E., Di Tizio, A., Marinari, S. (2010): Soil carbon dioxide emission and carbon content as affected by conventional and organic cropping systems in Mediterranean environment. – *Appl Soil Ecol* 46: 64-72.
- [26] McLean, E. O. (1982): Soil pH and Lime Requirement. – In: Page, A. L., Miller, R. H., Keeney, D. R. (eds.) *Methods of Soil Analysis, Part 2*. ASA, Madison, WI.
- [27] Morell, F. J., Alvaro-Fuentes, J., Lampurlanes, J., Cantero-Martinez, C. (2010): Soil CO₂ fluxes following tillage and rainfall events in a semiarid Mediterranean agroecosystem: effects of tillage systems and nitrogen fertilization. – *Agr Ecosyst Environ* 139: 167-173.

- [28] Morell, F. J., Whitmore, A. P., Álvaro-Fuentes, J., Lampurlanés, J., Cantero-Martínez, C. (2012): Root respiration of barley in a semiarid Mediterranean agroecosystem: field and modelling approaches. – *Plant Soil* 351: 135-147.
- [29] Meng-yang, Y., Ya-ru, Y., Lu-jun, L., Yan-Li, X., Xiao-zeng, H. (2014): Soil CO₂ emissions as affected by 20-year continuous cropping in Mollisols. – *Journal of Integrative Agriculture* 13: 615-623.
- [30] Nelson, D. W., Sommers, L. E. (1982): Chemical and Microbiological Properties. – In: Page, A. L., Miller, R. H. Keeney, D. R. (eds.) *Methods of Soil Analysis, Part 2*. ASA, Madison, WI.
- [31] Ngao, J., Epron D., Delpierre N., Bréda N., Granier A., Longdoz B. (2012): Spatial variability of soil CO₂ efflux linked to soil parameters and ecosystem characteristics in a temperate beech forest. – *Agricultural and Forest Meteorology* 154: 136-46.
- [32] Olsen, S. R., Sommers, L. E. (1982): Phosphorus. – In: Page, A. L., Miller, R. H., Keeney, D. R. (eds.) *Methods of Soil Analysis, Part 2*. ASA, Madison, WI.
- [33] Panosso, A. R., Marques, J. Jr., Pereira, G. T., La Scala, N. Jr. (2009): Spatial and temporal variability of soil CO₂ emission in a sugarcane area under green and slash-and-burn managements. – *Soil Till Res.* 105: 275-282.
- [34] Panosso, A. R., Marques Jr, J., Milori, D. M. B. P., Ferraudo, A. S., Barbieri, D. M., Pereira, G. T., La Scala Jr., N. (2011): Soil CO₂ emission and its relation to soil properties in sugarcane areas under slash-and-burn and green harvest. – *Soil and Tillage Research* 111: 190-196.
- [35] Raich, J. W., Potter, C. S., Bhagawati, D. (2002): Interannual variability in global soil respiration, 1980-94. – *Global Change Biol* 8: 800-812.
- [36] Rastogi, M., Singh, S., Pathak, H. (2002): Emission of carbon dioxide from soil. – *Current Science* 82: 510-517.
- [37] Reth, S., Reichstein, M., Falge, E. (2005): The effect of soil water content, soil temperature, soil pH-value and the root mass on soil CO₂ efflux—a modified model. – *Plant Soil* 268: 21-33.
- [38] Sauerbeck, D. R., (2001): CO₂ emissions and C sequestration by agriculture. Perspectives and limitations. – *Nutrient Cycling in Agroecosystems* 60: 253-266.
- [39] Xu, Q., Yang, R., Dong, Y.-X., Liu, Y.-X., Qiu, L.-R. (2016): The influence of rapid urbanization and land use changes on terrestrial carbon sources/sinks in Guangzhou, China. – *Ecological Indicators* 70: 304-316.
- [40] Wang, L. G., Qiu, J. J., Tang, H. J., Hu, L., Li, C. S., Eric, V. R. (2008): Modeling soil organic carbon dynamics in the major agricultural regions of China. – *Geoderma* 147: 47-55.
- [41] Zhang, H., Zhou, X., Lu, F., Pang, J., Feng, Z., Liu, W., O, Z., Wang, X. (2011): Seasonal dynamics of soil CO₂ efflux in a conventional tilled wheat field of the Loess Plateau, China. – *Ecol Res* 26: 735-743.
- [42] Zhang, T., Wang, G., Yang, Y., Mao, T., Chen, X. (2015): Non-growing season soil CO₂ flux and its contribution to annual soil CO₂ emissions in two typical grasslands in the permafrost region of the Qinghai-Tibet Plateau. – *European Journal of Soil Biology* 71: 45-52.