

EFFECTS OF EXOGENOUS CaCl₂ ON THE PHOTOSYNTHETIC FUNCTION AND ACTIVE OXYGEN METABOLISM OF *SALIX VIMINALIS* LEAVES UNDER PB STRESS

NING, W. – ZHU, S. Q. – NIAN, H. – YIN, J. H.* – AN, B. Y.*

College of Horticulture, Jilin Agricultural University, Changchun 130118, China

*Corresponding authors
e-mail:6325695@qq.com; jlnabya2020@126.com

(Received 9th Aug 2021; accepted 3rd Dec 2021)

Abstract. In order to provide some basic data for revealing the mechanism of exogenous CaCl₂ on improving drought resistance of *Salix viminalis* seedlings. The experimental results showed that: Spraying exogenous of different CaCl₂ concentrations significantly alleviated the damage degree of *S. viminalis* seedlings leaves caused by heavy metal stress, and the effect of 30 μmol·L⁻¹ CaCl₂ was the most significant. Spraying exogenous CaCl₂ could regulate stomatal limitation of *S. viminalis* seedlings leaves under heavy metal stress, which was beneficial to water holding capacity of *S. viminalis* seedlings leaves under heavy metal stress, and enhanced photosynthetic carbon assimilation capacity of leaves under heavy metal stress; Spraying exogenous CaCl₂ can reduce the energy pressure of PSII reaction center by increasing the non-photochemical quenching (NPQ) of leaves of *S. viminalis* seedlings, alleviate the photoinhibition of PSII, and promote the electron transfer process, especially on the receptor side of PSII; Spraying exogenous CaCl₂ effectively reduced the production of reactive oxygen species in leaves of *S. viminalis* seedlings, as well as the degree of membrane peroxidation, which was also one of the important reasons for alleviating the inhibition of photosynthetic capacity.

Keywords: heavy metals, *S. viminalis*, PSII, reactive oxygen, antioxidants

Introduction

Heavy metal stress is an important limiting factor in agricultural production, and the yield of crops caused by heavy metal stress alone exceeds the sum of all pathogens. Lead (Pb) is one of the most harmful heavy metals in the environment. In recent years, with the rapid development of economy, mining, agriculture and recycling of waste metals have significantly increased the concentration of Pb in the soil, and the damage dealt by Pb pollution to the environment has gradually intensified, and the impact on human health has become a hot topic (Soffianian et al., 2014; Sauliutė and Svecėvičius, 2015). The majority of Pb enters the human body through the soil-plant-human pathway and causes harm. To a certain extent, Pb in the soil will have a series of adverse effects on plant physiological metabolism, such as the increase of cell membrane permeability, the accumulation of reactive oxygen species, and the aggravation of membrane lipid peroxidation, and even lead to plant death in severe cases (Shu et al., 2012; Ogbomida et al., 2018).

Plant photosynthesis is one of the most sensitive processes to heavy metals. Persistent heavy metal stress will cause irreversible damage to plant photosynthetic apparatus (Zhang et al., 2018a), such as inhibition of photosynthetic pigment degradation (Albert et al., 2011; Guadagno et al., 2017; Chen et al., 2018; Zhang et al., 2018), photosynthetic phosphorylation and electron transport. Oxidative stress caused by oxidative stress in plant cell membrane caused by oxidative stress (Gill et al., 2010; Wang et al., 2017). Plant photosystem II (PSII) is one of the sensitive parts to heavy metal stress. A series of

photosynthetic physiological processes such as light energy absorption, water photolysis and electron transfer are closely related to PSII (Allahverdiyeva et al., 2013; Chen et al., 2017). To ensure the stability of photosynthetic function, especially PSII function, of plant leaves under heavy metal stress plays an important role in maintaining the normal growth of plants and improving the stress resistance of plants.

Calcium is not only an essential mineral nutrient element for plants, but also a second messenger for intracellular physiological and biochemical reactions. Therefore, the stress resistance of plants can be improved by stabilizing cell wall and cell membrane structures and inducing the expression of specific genes (Ryan and Kochian, 1993; Larkindale and Huang, 2004). At present, there are a lot of reports about CaCl₂ can protect the normal physiology of PSII and increase the content of antioxidant enzymes in plant leaves under stress. *S. viminalis* is a *Salix* plant of *Salix family*. *S. viminalis* has the advantages of rapid growth, high biomass and strong ability to accumulate heavy metals is widely used in phytoremediation and biomass energy development in heavy metal contaminated soil areas, and has certain ecological and economic value (Zhai et al., 2016). Therefore, improving the resistance of *S. viminalis* seedlings to heavy metal stress is the key to ensure the survival and growth of seedlings after transplanting. However, the regulation of exogenous CaCl₂ on physiological function of *S. viminalis* seedlings under heavy metal stress, especially the regulation mechanism of PSII function, was less studied. In this experiment, the effects of spraying different concentrations of exogenous CaCl₂ on PSII function of *S. viminalis* leaves under heavy metal stress were studied, in order to provide some basic data for improving the drought resistance ability of *S. viminalis*.

Materials and methods

Test materials and treatment

The experiment was conducted in the laboratory of soil science of Jilin Agricultural University (Changchun, Jilin Province, China) from March to June 2019. The annual seedlings were raised by cutting. The culture substrate was fully mixed with peat soil and quartz sand; the ratio was 2:1 (V / V). It was cultured in an artificial climate box with temperature of 25/23 °C (light / dark), light intensity of 400 μmol·m⁻²·s⁻¹, photoperiod of 12/12 h (light/dark), and relative humidity of about 75%. After the seedlings had long functional leaves, 1/2 of Hoagland nutrient solution was irrigated once a week. When the seedlings grow to six leaves and one heart, they are transplanted. Before transplanting, the seedlings are irrigated once to make the relative water content of the soil basically reach saturation. The seedlings are transplanted into a cultivation bowl with a diameter of 12 cm and a height of 15 cm, and one plant is planted in each pot.

After transplanting, the seedlings with the same growth were selected as the experimental seedlings. The treatment group was sprayed with CaCl₂ solution with concentration of 15 and 30 μmol·L⁻¹ (PH=7), and the control group was sprayed with water (CK). The foliar spray was carried out at 4:00 p.m., and the front and back sides were evenly sprayed until the solution on the leaves formed fine mist like uniform droplets ready to drop, after the water on the leaf surface evaporates naturally and the CaCl₂ solution is completely absorbed, repeat spraying 1 time with the same amount each time. 10 plants in each treatment were repeated. After spraying CaCl₂, the water on the surface of leaves is naturally evaporated. In order to fully absorb the spraying CaCl₂ on the leaf surface, after the CaCl₂ solution on the leaf surface evaporates, spray once more, and the usage and dosage are the same as the first time. The content of Pb in the test

matrix was set at 200 mg·kg⁻¹. On the 15th day after treatment, the physiological indexes were determined after the results showed obvious differences among different treatments.

Determination parameters and methods

The growth parameters were determined: Vernier caliper was used to measure plant height and leaf area was measured by leaf area meter.

Photosynthetic gas exchange parameters were measured: A CIRAS-2 portable photosynthesis system (PPsystem Company, UK) was used to determine the net photosynthetic rate (P_n), stomatal conductance (G_s), transpiration rate (T_r), and intercellular CO₂ concentration (C_i) of the second fully expanded functional leaf of the penultimate leaf of *S. viminalis* seedlings was selected. The CO₂ concentration was fixed at 400 μl·L⁻¹ in a CO₂ cylinder. The light intensity PFD was set to 1000 μmol·m⁻²·s⁻¹ with the light source built in the instrument. The net photosynthetic rate (P_n), stomatal conductance (G_s), transpiration rate (T_r) and intercellular CO₂ concentration (C_i) of *S. viminalis* leaves under different treatments were measured. Repeat 5 times.

Chlorophyll fluorescence parameters were measured: The electron transfer rate (ETR) and non-photochemical quenching (NPQ), the maximum photochemical efficiency (F_v/F_m) and the actual photochemical efficiency (Φ_{PSII}) of PSII reaction center under light acclimation were measured by FMS-2 (Hansatech company, UK) for 5 times.

The chlorophyll fluorescence kinetic curve and its parameters were determined: After 30 min dark adaptation, the OJIP curve of leaves after dark adaptation was measured by handy pea (Hansatech company, UK). According to Strasser et al.'s method (1995), the OJIP curves were standardized by $V_{O-P}=(F_t-F_o)/(F_m-F_o)$ and $V_{O-J}=(F_t-F_o)/(F_J-F_o)$ respectively. The relative variable fluorescence (V_J and V_K) of J point at 2 ms and K point at 0.3 ms were obtained, respectively. The differences between the standardized V_{O-P} and V_{O-J} curves of different treatments and the control were calculated, expressed as V_{O-P} and V_{O-J} , respectively. The measured OJIP curve was analyzed by JIP test. The maximum photochemical efficiency (F_v/F_m) of PSII and the photosynthetic performance index (PI_{ABS}) based on absorbed light energy were measured.

Determination of ROS metabolism and other physiological indices: The rate of production of O₂[·] was measured using the method of Zhang et al. (2007), The malondialdehyde (MDA) content was determined using the thiobarbituric acid method. The conductivity was measured by DDS-11C. Relative conductivity was used to express the electrolyte leakage rate. The content of superoxide dismutase activity (SOD) was determined using the NBT method. The activity unit (U) was 50% of the enzyme that inhibited the photochemical reduction of NBT in 1 ml of reaction solution in 1 h. The activity of ascorbic acid peroxidase (APX) was determined as described by Shen et al. (1996). An activity unit (U) is defined as the amount of enzyme that catalyzes 1 μmol ascorbic acid oxidation in one minute. Each index was measured five times.

Data and analysis

Excel and SPSS software (Version. 22) were used to conduct statistical analyses on the measured data. The data in the figure was denoted as mean ± standard deviation (SE). One-way ANOVA and least significant difference (LSD) were used to compare the differences among different data groups.

Results and analysis

Effects of exogenous CaCl₂ on leaf growth of S. viminalis seedlings under Pb Stress

Under Pb stress, the leaf growth of *S. viminalis* seedlings changed significantly, and the increase of leaf area of *S. viminalis* seedlings without spraying exogenous CaCl₂ was significantly delayed (Fig. 1). Compared with CK, the plant height of *S. viminalis* seedlings was decreased by 20.36% ($P < 0.01$), reaching a very significant difference level. However, spraying different concentrations of exogenous CaCl₂ significantly alleviated the reduction of leaf area of *S. viminalis* seedlings under heavy metal stress, of which 15 and 30 $\mu\text{mol}\cdot\text{L}^{-1}$ were sprayed under heavy metal stress. The leaf area of *S. viminalis* seedlings treated with 30 $\mu\text{mol}\cdot\text{L}^{-1}$ exogenous CaCl₂ was 26.51% ($P < 0.05$) and 37.21% ($P < 0.05$), respectively.




Figure 1. Effects of CaCl₂ on leaf growth of *S. viminalis* seedlings under Pb Stress. CK: CK + 0 CaCl₂ contents; +CaCl₂: CK + different amounts of CaCl₂ contents. Note: Data in the figure are the mean \pm SD; values followed by different lowercase letters indicate a significant difference ($p < 0.05$)

Effects of exogenous CaCl₂ on Photosynthetic gas exchange parameters of S. viminalis seedlings under Pb Stress

Compared with CK, P_n , G_s and T_r of *S. viminalis* seedlings leaves under Pb stress were significantly decreased, but spraying different concentrations of exogenous CaCl₂ significantly alleviated the reduction range of P_n , G_s and T_r . Except that there was no significant difference between spraying different concentrations of exogenous CaCl₂ and spraying CaCl₂ under heavy metal stress, spraying 15 and 30 $\mu\text{mol}\cdot\text{L}^{-1}$ in leaves of *S. viminalis* seedlings leaves Under CaCl₂ Treatment, P_n and G_s in leaves of *S. viminalis* seedlings were significantly increased compared with those without CaCl₂ treatment. It can be seen from Fig. 2D that spraying different concentrations of exogenous CaCl₂ also significantly alleviated the C_i reduction of *S. viminalis* seedlings under heavy metal stress, but there was no significant difference between the C_i of *S. viminalis* seedling leaves treated with 30 $\mu\text{mol}\cdot\text{L}^{-1}$ exogenous CaCl₂ and that without spraying exogenous CaCl₂.

Effects of exogenous CaCl₂ on F_v/F_m , Φ_{PSII} , ETR and NPQ of S. viminalis seedlings under Pb stress

Under Pb stress, F_v/F_m , Φ_{PSII} and ETR of *S. viminalis* seedlings decreased significantly, while NPQ increased significantly (Fig. 3). However, spraying different concentrations of CaCl₂ significantly alleviated the decrease of F_v/F_m , Φ_{PSII} and ETR of *S. viminalis* seedlings leaves, especially under heavy metal stress, when spraying

30 μmol·L⁻¹ exogenous CaCl₂, the Φ_{PSII} of *S. viminalis* seedlings leaves increased by 73.24% (*P*<0.01). There was no significant difference in *F_v/F_m*, Φ_{PSII} and ETR between different concentrations of CaCl₂. Spraying different concentrations of CaCl₂ increased NPQ of *S. viminalis* seedlings leaves under heavy metal stress, but there was no significant difference among different concentrations of CaCl₂. However, NPQ of *S. viminalis* seedlings under 30 μmol·L⁻¹ CaCl₂ treatment was significantly higher than that under Pb stress.




Figure 2. Effects of different concentrations of exogenous CaCl₂ on net photosynthetic rate (a), stomatal conductance (b), transpiration rate (c) and intercellular CO₂ concentration (d) of *S. viminalis* seedlings under Pb Stress. CK: CK + 0 CaCl₂ contents; +CaCl₂: CK + different amounts of CaCl₂ contents. Note: Data in the figure are the mean ± SD values followed by different lowercase letters indicate a significant difference (*p*<0.05)

Effects of exogenous CaCl₂ on OJIP curve of *S. viminalis* seedlings under Pb stress

Compared with CK, the relative fluorescence intensity of O point and J point had no significant change under Pb stress, while the relative fluorescence intensity of I and P decreased significantly, especially that of P point (Fig. 4). However, spraying different concentrations of exogenous CaCl₂ could significantly reduce the change range of OJIP curve of *S. viminalis* seedlings under heavy metal stress.

Effects of exogenous CaCl₂ on standardized O-P curve, *V_J* and *V_I* of *S. viminalis* seedlings under heavy metal stress

Compared with CK, the relative variable fluorescence (*V_J*) of each point on the *V_{O-P}* curve of *S. viminalis* seedlings leaves under Pb stress increased significantly with the relative variable fluorescence *V_J* of J point at 2 ms (Fig. 5). the increase range of *V_J* in leaves of *S. viminalis* seedlings treated with different concentrations of exogenous CaCl₂ was significantly less than that of no CaCl₂ treatment. At 0.3 ms, the change range of K point relative to variable fluorescence *V_K* was small, and it was not significantly affected by spraying CaCl₂.




Figure 3. Effects of different concentrations of exogenous CaCl₂ on the maximum photochemical efficiency (F_v/F_m) and the actual photochemical efficiency (Φ_{PSII}) of PSII reaction center electron transfer rate (ETR) and non-photochemical quenching (NPQ) in leaves of *S. viminalis* seedlings under Pb Stress. CK: CK + 0 CaCl₂ contents; +CaCl₂: CK + different amounts of CaCl₂ contents. Note: Data in the figure are the mean \pm SD values followed by different lowercase letters indicate a significant difference ($p < 0.05$)




Figure 4. Effects of different concentrations of exogenous CaCl₂ on OJIP curve of *S. viminalis* seedlings under Pb Stress. CK: CK + 0 CaCl₂ contents; +CaCl₂: CK + different amounts of CaCl₂ contents

The results of quantitative analysis of V_J and V_K showed that the V_J of *S. viminalis* seedling leaves under Pb stress increased by 24.63% ($P < 0.05$), while the V_J of leaves of *S. viminalis* seedlings treated with 15 and 30 $\mu\text{mol}\cdot\text{L}^{-1}$ CaCl₂ increased by 21.77% ($P < 0.05$) and 18.44% ($P < 0.05$), respectively. Under Pb stress, the V_K of *S. viminalis* seedlings leaves had no significant change compared with CK, and spraying different concentrations of CaCl₂ did not significantly change the V_K of *S. viminalis* seedlings leaves.




Figure 5. Effects of different concentrations of exogenous CaCl₂ on V_j (a) and V_c (b) in leaves of *S. viminalis* seedlings under Pb Stress. CK: CK + 0 CaCl₂ contents; +CaCl₂: CK + different amounts of CaCl₂ contents. Note: Data in the figure are the mean ± SD; values followed by different lowercase letters indicate a significant difference ($p < 0.05$)

Effects of exogenous CaCl₂ on reactive oxygen species and membrane peroxidation of *S. viminalis* seedlings under Pb Stress

Compared with CK, the results showed that the treatment of 30 µmol·L⁻¹ exogenous CaCl₂ had the most obvious effect, and the O₂^{·-} production rate and H₂O₂ content decreased by 25.83% ($P < 0.05$) and 32.22% ($P < 0.05$), respectively, which resulted in the decrease of MDA content and electrolyte leakage rate of membrane lipid peroxidation by 21.43% ($P < 0.05$) and 25.14% ($P < 0.05$), respectively (Fig. 6).




Figure 6. Effects of different concentrations of exogenous CaCl₂ on O₂^{·-} production rate (a), H₂O₂ Content (b), MDA content (c) and electrolyte leakage rate (d) in leaves of *S. viminalis* seedlings under Pb Stress. CK: CK + 0 CaCl₂ contents; +CaCl₂: CK + different amounts of CaCl₂ contents. Note: Data in the figure are the mean ± SD; values followed by different lowercase letters indicate a significant difference ($p < 0.05$)

Discussion

Pb is a kind of non-essential element for plant growth, which has different effects on different plants when it enters into plants. Many studies have shown that low concentrations of Pb can stimulate plant growth, moderate concentrations of Pb can cause plant metabolic disorders and inhibit growth, while heavy concentrations of Pb can cause plant death (Del et al., 1999a,b; Dong et al., 2008), CaCl₂ could alleviate the growth inhibition of heavy metals on plants (Saradhi et al., 2001; Li et al., 2011; Zhang et al., 2014). In this study, Pb significantly reduced the growth rate of plant height and leaf area; Application of CaCl₂ after Pb stress could reduce the inhibition of Pb on leaf area and plant height growth of *S. viminalis*.

Under Pb stress, the photosynthetic carbon assimilation capacity of *S. viminalis* seedlings leaves was limited, which was mainly manifested in the decrease of P_n , along with the decrease of G_s , T_r and C_i . According to Farquhar's photosynthetic stomatal factor analysis theory (Farquhar et al., 2003), the reason for the decrease of photosynthetic carbon assimilation capacity of *S. viminalis* seedlings under heavy metal stress was directly related to the decrease of stomatal conductance. Although the decrease of water loss can effectively prevent the loss of water, it also directly reduces the supply of carbon assimilation material (CO₂), which limits its carbon assimilation capacity. In this experiment, spraying different concentrations of exogenous CaCl₂ increased P_n in different degrees, which was similar to the change of G_s , and accompanied by the increase of C_i , indicating that spraying exogenous CaCl₂ could improve the photosynthetic capacity of *S. viminalis* seedlings by improving stomatal limitation.

Under Pb stress, the dark response of plant leaves was inhibited, and the accumulation of assimilative capacity (ATP and NADPH) would feedback inhibit the light response process, resulting in the excess of electrons in the photosynthetic electron transport chain (Li et al., 2000; Liu et al., 2006). In addition, the decrease of PSII activity also inhibited the process of light energy absorption and electron transfer. In this experiment, although F_v/F_m , ETR and, Φ_{PSII} of *S. viminalis* seedlings were significantly decreased under Pb stress. F_v/F_m and, Φ_{PSII} were important indexes reflecting the photochemical activity of PSII, and the sensitivity of Φ_{PSII} was significantly higher than that of F_v/F_m (Kalaji et al., 2016). Therefore, the results showed that the photochemical activity of PSII in leaves of *S. viminalis* seedlings was inhibited under heavy metal stress, and the process of PSII electron transport was hindered. However, the decrease of PSII photochemical activity was alleviated by spraying different concentrations of exogenous CaCl₂, especially 30 $\mu\text{mol}\cdot\text{L}^{-1}$. The results showed that exogenous CaCl₂ could alleviate the photoinhibition of *S. viminalis* seedlings under heavy metal stress and promote electron transfer. NPQ was positively correlated with heat dissipation dependent on xanthophyll cycle. Under Pb stress, NPQ in leaves of *S. viminalis* seedlings increased significantly, which means that the excess excitation energy in PSII could be dissipated by increasing NPQ under Pb stress, so as to reduce the pressure of PSII reaction center. The NPQ of *S. viminalis* seedlings leaves treated with different concentrations of exogenous CaCl₂ increased to varying degrees. Ivanov et al. (1995) found that CaCl₂ treatment enhanced the xanthophyll cycle in barley seedlings, which enhanced the opening degree of photosystem II reaction center and promoted the utilization rate of light energy. Therefore, the reason why spraying exogenous CaCl₂ to alleviate the decrease of PSII in leaves of *S. viminalis* seedlings under heavy metal stress may be related to the mechanism of energy dissipation dependent on xanthophyll cycle induced by CaCl₂.

Under stress conditions, the blocking sites of photosynthetic electron transfer often occur on the electron donor side and receptor side of PSII reaction center, especially the transfer of Q_A to Q_B is the main inhibition site (Zhang et al., 2018a). The increase of J point relative to variable fluorescence V_J at 2 ms on the normalized O-P curve indicated that the electron transfer from Q_A to Q_B was blocked in the photosynthetic electron transport chain (Zhang et al., 2017; Zhang, 2018b). The increase of relative variable fluorescence V_K at 0.3 ms on the normalized O-J curve was considered as a specific marker of damage to OEC activity of PSII electron donor side oxygen complex (Zhang et al., 2012, 2016). In this experiment, the V_J of *S. viminalis* seedlings leaves increased significantly under Pb stress, but the V_K did not change significantly, which indicated that the reason for the decrease of PSII photosynthetic electron transfer rate under Pb stress mainly occurred in PSII receptor side. However, the change of V_K was not only affected by the donor side injury of PSII, but also by the receptor side injury of PSII. When the injury degree of the recipient side was greater than that of the donor side, the V_K did not increase significantly (Xu et al., 2018; Zhang et al., 2018c). Therefore, in this study, the effect of heavy metal stress on the donor side of PSII may be due to the insensitivity of OEC to Pb stress, or it may be due to the excessive damage on the receptor side. However, spraying different concentrations of exogenous CaCl₂ significantly alleviated the increase of V_J, but the effect on V_K was not significant, which indicated that exogenous CaCl₂ could promote the electron transfer from Q_A to Q_B in PSII receptor side of *S. viminalis* seedlings under Pb stress.

Pb usually leads to excessive reduction of photosynthetic electron transport chain in plants, and produces a large number of ROS in chloroplasts and mitochondria (Ramachandra et al., 2004; Asada, 2009; Ahmed et al., 2009). Excessive ROS breaks the redox balance in plants, causes membrane peroxidation, damages membrane system, and causes oxidative damage to cell components and structures (Jiang and Zhang, 2004; Gill et al., 2010). Under normal conditions, the production and elimination of intracellular free radicals are in a dynamic equilibrium state. When the concentration of exogenous Pb²⁺ reaches a certain level, this dynamic balance will be broken, resulting in membrane lipid peroxidation and osmotic stress (Amhed et al., 2009; Li and Li, 2011). The results also showed that under Pb stress, the relative permeability of cell membrane increased, the production rate of O₂^{•-} and the content of MDA increased, which caused lipid peroxidation in the inner membrane of *S. viminalis*. Under Pb stress, O₂^{•-} production rate and H₂O₂ content were significantly increased, MDA content and electrolyte leakage rate were also significantly increased, which indicated that excessive ROS caused membrane system peroxidation and increased membrane permeability. However, spraying different concentrations of exogenous CaCl₂ significantly alleviated ROS production and membrane lipid peroxidation of *S. viminalis* seedlings, especially the treatment of 30 μmol·L⁻¹ CaCl₂ was the most effective, which was consistent with the change of photosynthetic parameters of *S. viminalis* seedlings. Spraying exogenous CaCl₂ could alleviate ROS production by enhancing photosynthetic capacity and reducing PSII damage degree of *S. viminalis* seedlings, and the decrease of ROS production could also alleviate the photoinhibition of *S. viminalis* seedlings leaves under heavy metal stress.

Conclusion

The photosynthetic capacity of *S. viminalis* seedlings was decreased under Pb stress, mainly due to stomatal factors, and also related to the decrease of PSII photochemical

activity and the damage of membrane system caused by ROS. Under Pb stress, spraying different concentrations (15 and 30 $\mu\text{mol}\cdot\text{L}^{-1}$) of exogenous CaCl₂ promoted the photosynthetic capacity of *S. viminalis* seedlings, especially 30 $\mu\text{mol}\cdot\text{L}^{-1}$ CaCl₂. The main reason is that exogenous CaCl₂ can effectively reduce the energy pressure of PSII reaction center in leaves of *S. viminalis* seedlings under Pb stress by inducing the increase of NPQ, thus alleviating the photoinhibition degree of PSII under Pb stress, promoting the electron transfer rate, especially the electron transfer ability of PSII receptor side. Under Pb stress, spraying exogenous CaCl₂ could also effectively reduce ROS production in leaves of *S. viminalis* seedlings and alleviate the degree of membrane lipid peroxidation. Therefore, the inhibition of ROS production by spraying CaCl₂ under heavy metal stress is not only related to the enhancement of photosynthetic capacity, but also to the increase of enzymatic or non-enzymatic related ROS scavenging system induced by CaCl₂, but this needs further study.

Funding. This study was financially supported by Jilin Provincial Natural Science Foundation No. YDZJ202101ZYTS113.

REFERENCES

- [1] Ahmed, C. B., Rouina, B. B., Sensoy, S., Boukhris, M., Abdallah, F. B. (2009): Changes in gas exchange, proline accumulation and antioxidative enzyme activities in three olive cultivars under contrasting water availability regimes. – *Environmental & Experimental Botany* 67(2): 345-352.
- [2] Albert, K. R., Mikkelsen, T. N., Michelsen, A., Ropoulsen, H., Leon, V. D. L. (2011): Interactive effects of drought, elevated CO₂ and warming on photosynthetic capacity and photosystem performance in temperate heath plants. – *Journal of Plant Physiology* 168(13): 1550-1561.
- [3] Allahverdiyeva, Y., Suorsa, M., Rossi, F., Pavesi, A., Kater, M. M., Antonacci, A., Tadini, L., Pribil, M., Schneider, A., Wanner, G., Leister, D., Aro, E. M., Barbato, R., Pesaresi, P. (2013): Arabidopsis plants lacking PsbQ and PsbR subunits of the oxygen-evolving complex show altered PSII supercomplex organization and short-term adaptive mechanisms. – *Plant Journal* 75(4): 671-684.
- [4] Asada, K. (2006): Production and scavenging of reactive oxygen species in chloroplasts and their functions. – *Plant Physiology* 141(2): 391-396.
- [5] Chen, E., Zhang, C. M., Su, Y. Q., Ma, J., Zhang, Z. W., Yuan, M., Zhang, H. Y., Yuan, S. (2017): Responses of photosystem II and antioxidative systems to high light and high temperature co-stress in wheat. – *Environmental & Experimental Botany* 135(3): 45-55.
- [6] Chen, Z. F., Wang, Z., Yang, Y. G., Li, M., Xu, B. C. (2018): Abscisic acid and brassinolide combined application synergistically enhances drought tolerance and photosynthesis of tall fescue under water stress. – *Scientia Horticulturae* 228(26): 1-9.
- [7] Del, V. C., Barea, J. M., Azcon-Aguilar, C. (1999a): Diversity of arbuscularmycorrhizal fungus populations in heavy-metal-contaminated soils. – *Applied Environmental Microbiology* 65(2): 718-723.
- [8] Del, V. C., Barea, J. M., Azcon-Aguilar, C. (1999b): Assessing the tolerance to heavy metals of arbuscular mycorrhizal fungi isolated from sewage sludge-contaminated soils. – *Applied Soil Ecology* 11(2-3): 261-269.
- [9] Dong, Y., Zhu, Y. G., Smith, F. A., Wang, Y., Chen, B. (2008): Arbuscular mycorrhiza enhanced arsenic resistance of both white clover (*Trifolium repens* Linn.) and ryegrass (*Lolium perenne* L.) plants in an arsenic-contaminated soil. – *Environmental Pollution* 155(1): 174-181.

- [10] Farquhar, G. D. F., Sharkey, T. D. (2003): Stomatal Conductance and Photosynthesis. – Annual Review of Plant Physiology and Plant Molecular Biology 33(33): 317-345.
- [11] Gill, S. S., Tuteja, N. (2010): Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. – Plant Physiology & Biochemistry 48(12): 909-930.
- [12] Guadagno, C. R., Ewers, B. E., Speckman, H. N., Aston, T. L., Huhn, B. J., DeVore, S. B., Ladwig, J. T., Strawn, R. N., Weinig, C. (2017): Dead or alive? Using membrane failure and chlorophyll fluorescence to predict mortality from drought. – Plant Physiology 175(1): 223-234.
- [13] Hu, Y. B., Sun, G. Y., Wang, X. C. (2007): Induction characteristics and response of photosynthetic quantum conversion to changes in irradiance in mulberry plants. – Journal of Plant Physiology 164(8): 959-968.
- [14] Ivanov, A. G., Krol, M. D., Huner, N. P. (1995): Abscisic acid induced protection against photoinhibition of PSII correlates with enhanced activity of the xanthophyll cycle. – Febs Letters 371(1): 61-64.
- [15] Jiang, M. Y., Zhang, J. H. (2004): Abscisic acid and antioxidant defense in plant cells. – Acta Botanica Sinica 46(1): 1-9.
- [16] Kalaji, H. M., Schansker, G., Ladle, R. J., Goltsev, V., Bosa, K., Allakhverdiev, S. I., Brestic, M., Bussotti, F., Calatayud, A., Dabrowski, P., Elsheery, N. I., Ferroni, L., Guidi, L., Hogewoning, S. W., Jajoo, A., Misra, A. N., Nebauer, S. G., Pancaldi, S., Penella, C., Poli, D., Pollastrini, M., Romanowska-Duda, Z. B., Rutkowska, B., Serodio, J., Suresh, K., Szulc, W., Tambussi, E., Yannicari, M., Zivcak, M. (2014): Frequently asked questions about in vivo chlorophyll fluorescence: practical issues. – Photosynthesis Research 122(2): 121-58.
- [17] Larkindale, J., Huang, B. R. (2004): Thermotolerance and antioxidant systems in *Agrostis stolonifera*: involvement of salicylic acid, abscisic acid, calcium, hydrogen peroxide, and ethylene. – Journal of Plant Physiology 161(4): 405-413.
- [18] Li, X. P., Bjorkman, O., Shi, C., Grossman, A. R., Rosenquist, M., Jansson, S., Niyogi, K. K. (2000): A pigment-binding protein essential for regulation of photosynthetic light harvesting. – Nature 403(6768): 391-395.
- [19] Li, Y. Q., Li, T. L. (2011): Regulations of Calcium and Calcium Inhibitors on the Low Light Tolerance in Tomato. – Acta Agriculturae Boreali-Occidentalis Sinica.
- [20] Liu, W. J., Yuan, S. N. H., Lei, T., Duan, H. G., Liang, H. G., Lin, H. H. (2006): Effect of water stress on photosystem 2 in two wheat cultivars. – Biologia Plantarum 50(4): 597-602.
- [21] Ogbomida, E. T., Nakayama, S. M. M., Bortey-Sam, N., Oroszlany, B., Tongo, I., Enuneku, A. A., Ozekeke, O., Ainerua, M. O., Fasipe, I. P., Ezemonye, L. I., Mizukawa, H., Ikenaka, Y., Ishizuka, M. (2018): Accumulation patterns and risk assessment of metals and metalloids in muscle and offal of free-range chickens, cattle and goat in Benin City. – Ecotoxicology and Environmental Safety 151: 98-108.
- [22] Ramachandra, R. A., Chaitanya, K. V., Vivekanandan, M. (2004): Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. – Journal of Plant Physiology 161(11): 1189-1202.
- [23] Ryan, P. R., Kochian, L. V. (1993): Interaction between aluminum toxicity and calcium uptake at the root apex in near-isogenic lines of wheat (*Triticum aestivum* L.) differing in aluminum tolerance. – Plant Physiology 102(3): 975-982.
- [24] Saradhi, P. P., Suzuki, I., Katoh, A., Sakamoto, A., Sharmila, P., Shi, D. J. (2000): Protection against the photo-induced inactivation of the photosystemII complex by CaCl₂ cistic acid. – Plant Cell Environment 23(7): 711-718.
- [25] Sauliutė, G., Svecevičius, G. (2015): Heavy metal interactions during accumulation via direct route in fish: a review. – Zoology & Ecology 25(1): 77-86.
- [26] Shen, W. B., Xu, L. L., Ye, M. B., Zhang, R. X. (1996): Study on determination of ASP activity. – Plant Physiology Communications 32(3): 203-205.

- [27] Shu, X., Yin, L. Y., Zhang, Q. F., Wang, W. B. (2012): Effect of Pb toxicity on leaf growth, antioxidant enzyme activities, and photosynthesis in cuttings and seedlings of *Jatropha curcas* L. – Environmental Science and Pollution Research 19(3): 893-902.
- [28] Soffianian, A., Madani, E., Arabi, M. (2014): Risk assessment of heavy metal soil pollution through principal components analysis and false color composition in Hamadan Province, Iran. – Environmental Systems Research 3(1): 1-14.
- [29] Strasser, R. J., Srivastava, A., Govindjee (1995): Polyphasic chlorophyll a fluorescence transient in plants and cyanobacteria. – Photochemistry and Photobiology 61(1): 32-42.
- [30] Wang, Y. G., Peng, C. X., Zhan, Y. N., Yu, L. H., Li, M., Li, J., Geng, G. (2017): Comparative proteomic analysis of two sugar beet cultivars with contrasting drought tolerance. – Journal of Plant Growth Regulation 36(3): 1-13.
- [31] Xu, N., Zhang, H. H., Zhong, H. X., Wu, Y. N., Li, J. B., Li, X., Yin, Z. P., Zhu, W. X., Qu, Y., Sun, G. Y. (2018): The response of photosynthetic functions of F1 cutting seedlings from *Physocarpus amurensis* Maxim (♀) × *Physocarpus opulifolius* “Diabolo” (♂) and the parental leaves to salt stress. – Frontiers in Plant Science, DOI: 10.3389/fpls.00714.
- [32] Zhai, F. F., Mao, J. M., Liu, J. X., Peng, X., Han, L., Sun, Z. Z. (2016): Male and female subpopulations of *S. viminalis* present high genetic diversity and high long-term migration rates between them. – Frontiers in Plant Science 7: 330.
- [33] Zhang, F. Q., Wang, Y. S., Luo, Z. P., Dong, J. D. (2007): Effect of heavy metal stress on antioxidative enzymes and lipid peroxidation in leaves and roots of two mangrove plant seedlings (*Kandeliacandel* and *Bruguieragymnorrhiza*). – Chemosphere 67(1): 44-50.
- [34] Zhang, Z., Li, G., Gao, H., Zhang, L. T., Yang, C., Liu, P., Meng, Q. W. (2012): Characterization of photosynthetic performance during senescence in stay-green and quick-leaf-senescence *Zea mays* L. inbred lines. – Plos One 7(8): e42936.
- [35] Zhang, G. H., Liu, Y. F., Ni, Y., Meng, Z. J., Lu, T. (2014): Exogenous Calcium Alleviates Low Night Temperature Stress on the Photosynthetic Apparatus of Tomato Leaves. – Plos One 9(5): e97322.
- [36] Zhang, H. H., Zhong, H. X., Wang, J. F., Xu, N. (2016): Adaptive changes in chlorophyll content and photosynthetic features to low light in *Physocarpus amurensis* Maxim and *Physocarpus opulifolius* “Diabolo”. – Peer J 4(3): e2125.
- [37] Zhang, H. H., Xu, N., Li, X., Jin, W. W., Tian, Q., Sun, G. Y., Gu, S. Y. (2017): Overexpression of 2-Cys Prx increased salt tolerance of photosystem II in tobacco. – International Journal of Agriculture and Biology 19(4): 735-745.
- [38] Zhang, H. H., Xu, N., Sui, X., Long, J. H., Wu, Y. N., Li, J. B., Wang, J. F., Qu, Y., Sun, G. Y. (2018a): Arbuscular mycorrhizal fungi (*Glomus mosseae*) improves growth, photosynthesis and protects photosystem II in leaves of *Lolium perenne* L. under cadmium contaminated soil. – Frontiers in Plant Science, DOI: 10.3389/fpls.01156.
- [39] Zhang, H. H., Feng, P., Yang, W., Sui, X., Li, W., Zhang, R. T., Gu, S. Y., Xu, N. (2018b): Effects of flooding stress on the photosynthetic apparatus of leaves of two *Physocarpus* cultivars. – Journal of Forest Research 39(4): 1049-1059.
- [40] Zhang, H. H., Xu, N., Sui, X., Yin, Z. P., Zhu, W. X., Li, X., Sun, G. Y. (2018c): Photosystem II function response to drought stress in leaves of two alfalfa (*Medicago sativa*) varieties. – International Journal of Agriculture and Biology 20(5): 1012-1020.