

PHYTOCHEMICAL SCREENING AND BIOLOGICAL EFFECTS OF *LAURUS NOBILIS* (LAURACEAE) ESSENTIAL OIL AGAINST MOSQUITO LARVAE, *CULEX PIPIENS* (LINNAEUS, 1758) (DIPTERA: CULICIDAE) SPECIES

AISSAOUI, L.¹ – BOUAZIZ, A.² – BOUDJELIDA, H.^{3*} – NAZLI, A.⁴

¹Research Laboratory of Improvement and Development of Animal and Plant Production, Department of Biology and Animal Physiology, Ferhat Abbas University, Setif 19000, Algeria
(e-mail: linda.ask12@gmail.com)

²Laboratory of Terrestrial and Aquatic Ecosystems, Department of Biology, Mohamed Cherif Messaadia University, Souk-Ahras 41000, Algeria
(e-mail: a.bouaziz@univ-soukhras.dz)

³Laboratory of Applied Animal Biology, Department of Biology, Faculty of Sciences, Badji Mokhtar University, Annaba 23000, Algeria

⁴Department of Molecular Biology and Genetics, Istanbul University, Turkey
(e-mail: narda@stanbul.edu.tr)

*Corresponding author
e-mail: boudjelidahamid@yahoo.fr

(Received 5th Jun 2022; accepted 11th Nov 2022)

Abstract. The present study aimed to determine the phytochemical compounds and to evaluate the activities of the essential oil (EO) of *Laurus nobilis* against mosquito larvae, *Culex pipiens*. The EO was extracted by hydro-distillation from *L. nobilis* leaves and subsequently, their chemical profile was identified by gas chromatography coupled with mass spectrometry (GC-MS) method. 39 compounds, representing 99.98% of the crude oil were identified; namely 1.8-cineole (22.42%), isolongifolene (10.22%), 3-carene (7.74%), Alpha-zingiberene (6.69%), Eremophilene (4.87%), Aristola-1 (10), 8-diene (3.09%) icyclo, sabinene (4.13%), α -pinene (1.94%) and Beta-caryophyllene (0.36%). Overall, 21 of these components, out of 39, were over 1%. The *L. nobilis* EO toxicity was tested at different concentrations, ranging between 5 and 35 μ l/l, against third and fourth-instar of *Cx. pipiens* larvae. Mortality was recorded for both stages and their lethal concentrations (LC₅₀ and LC₉₀) were estimated at 3.74 μ l/l and 14.47 μ l/l for the third-instar larvae and 18 μ l/l and 39.08 μ l/l for the fourth-instar larvae, respectively. The bioassay results showed a larvicidal effect with a concentration response-relationship. Furthermore, the compound effects, applied at LC₅₀ and LC₉₀ against the fourth-instars larvae showed an increase in the development duration and a disturbance in reproduction. The investigation confirmed the toxicological effects of *L. nobilis* EO against mosquito larvae; consequently, it could be considered as a potent source for producing natural larvicidal agents and bio-insecticides for pest and insect vector control.

Keywords: toxicity, plant essential oil, larvicidal activity, medicinal plants, arthropods

Introduction

Mosquitoes (Diptera: Culicidae) are the most widely distributed arthropods in the world and, in addition to causing a great annoyance, are considered the most significant arthropod vectors of diseases. The role of several mosquito species of the Culicidae family, in bites and pruritus of humans has doubled their medical importance (Salavati et al., 2021). Furthermore, the involvement of these vectors in the transmission of some

pathogens among domesticated and wild birds and animals has put this family at the forefront of important medical and veterinary insects. It is known that several mosquito species belonging to the genera *Anopheles*, *Culex* and *Aedes* are the main vectors for pathogens of various diseases causing serious health problems with increasing mortality rate to human beings (Raja et al., 2018; Mouhamadou et al., 2020). *Culex pipiens* is recognized as the primary vector especially of arboviruses (arthropod-borne viruses), such as St. Louis Encephalitis (SLE), and it was the main force in spreading the West Nile virus (Lounibos, 2002). The domestic mosquito, *Cx. pipiens* is considered among the most abundant species in Algeria (Aïssaoui and Boudjelida, 2017). It is widely distributed in the urban and suburban areas, due to the presence of many artificial breeding sites practically throughout the year (Arroussi et al., 2021).

Pest populations control around the world is essentially dependent upon chemical pesticides and fumigants. Repeated uses of these expensive synthetic products have disrupted natural biological control systems and led to the resurgence of these pests, the development of insect resistance, and undesirable effects on the environment, non-target organisms, and human health (Minetti et al., 2020; Richards et al., 2020; Smith and Perfetti, 2020). Also, the increasing concern; over the pesticide residue level in different environmental sites and especially in food has encouraged the scientists to propose a new alternatives of conventional pesticides (Boudjelida et al., 2005). Although there has been much research on the potential of natural plants to control insect pests, the interest in herbal based products was subsequently reduced due to the advent of synthetic chemicals or insecticides. However, the interest in anti-mosquito products derived from natural origin is being revived with considerable attention because the continued applications of synthetic compounds have some drawbacks, including the widespread development of insecticide resistance among vector population and biological magnification through food chain (Rawani et al., 2010; Flores and O'Neill., 2018; Wilson et al., 2020). Scientists, whose investigating in the control of vector-borne diseases, thought mosquitoes can be easily targeted in their larvae stages, since they breed in water (Bouaziz et al., 2017; Djeddar et al., 2021; Dar and Jamal, 2021). Global scientific efforts were initiated to discover and to develop safe, effective, and sustainable alternatives to synthetic chemical insecticides (Muturi et al., 2017). Phytochemicals have a promising major role in mosquito control programs owed to their efficiency, biodegradability and lack of secondary effects (Amalraj et al., 2000; Gunasekaran et al., 2004).

In the last decades, human has extensively used various medicinal plant extracts, in particular essential oils, that therapeutic properties have tangled global attention (Dadalioglu and Evrendilek, 2004). Furthermore, the larvicidal activity of plant essential oils against mosquitoes has been tested and showed toxic effects (Govindarajan et al., 2012). Many studies, using plant extracts (Nabti and Bounechada, 2019; da Costa et al., 2020), in order to explain their mode of action, have reported physiological changes that occurred in neuro-endocrine system function controlling insect behaviour, growth, molting process and metamorphosis (Djeghader et al., 2018; Corzo et al., 2020).

The aromatic plant *Laurus nobilis* belongs to the Laurel family, Lauraceae. It is one of the evergreen trees, cultivated in many warm regions of the world and its native territories are the southern Mediterranean countries (Marzouki et al., 2011). The bioactive plant ingredient(s) can be obtained from the whole plant or from a specific part. The leaf essential oil of the woody plant *L. nobilis* (Derwich et al., 2009), has been

traditionally used as an effective remedy to treat some health problems related to epigastric bloating, flatulence and fungal diseases (Patrakar et al., 2012; Caputo et al., 2020). In Algeria, it is cultivated in different regions and its leaves and fruits are used in food, cosmetics and pharmaceutical industries. The plant is widely used by local people as a source of spice and for its medicinal properties. In addition, laurel essential oil is currently used in folk medicines for the treatment of different health problems, such as rheumatism and dermatitis (Kilic et al., 2004; Georgiev and Stoyanova, 2006). Its name is dedicated to Apollo, the ancient Greek sun god, that symbolizes peace and victory (Dmitry and Naida, 2019).

The purpose of the present study was to determine the chemical composition of essential oil of the local plant, *L. nobilis*. and to examine its toxicity against larval stages of *Cx. pipiens* mosquito and to evaluate some developmental and physiological effects in order to understand its mode of action.

Materials and methods

Rearing of mosquito larvae of Culex pipiens

The toxicological essays were conducted in the laboratory following the standard methods for testing larval susceptibility, according to World Health Organization (WHO, 1991). The mosquito larvae of *Cx. pipiens* were obtained from the laboratory colonies. They are reared under controlled conditions (temperature at 25 ± 2 °C and photoperiod cycle of 12 light/12 dark). Mosquito larvae are reared in Pyrex storage jars containing 1000 ml of stored tap water. Larvae were daily fed with fresh food composed of a mixture of Biscuit-dried yeast (75:25 by weight), till the pupal stage was reached. Subsequently, the resulting pupae were transferred, with the help of a dipper, to jars containing water and placed in a cubic cage (30 × 30 × 30 cm) covered with a net until adult emergence. The adults were fed with 10% sugar solution for a period of three days before they were provided by mobilized shaved mouse for blood feeding.

Plant material and essential oil extraction

Fresh aerial parts of *L. nobilis* were randomly harvested, during the study period from a high-plains region (Setif, Northeast Algeria; 36°03'N 5°31'E) and were transported to the laboratory for chemical analysis. After collection, the plant leaves were initially rinsed with distilled water, soaked in paper towel and air-dried for 24 h. A sample of 100 g of plant powder was hydrodistilled for 3 h using a Clevenger-type apparatus (Ildam, Ankara), and the obtained essential oil was stored at 4 °C in an amber bottle. The essential oil yield was estimated according to dry leaves mass which is estimated by the ratio between the weight of extracted oil and that of the treated plant, that is expressed as a percentage; using the following equation: $Y (\%) = (m/MO) \times 100$, where m represents the essential oil mass (g), MO represents the dry leaves mass (g) and Y is the essential oil yield (%).

Gas chromatography-mass spectrometry analysis

The plant *L. nobilis* EO was analyzed by gas chromatography-mass spectrophotometry (GC-MS) equipped with a 5% phenyl-methylpolysiloxane DB-5MS column (30 m length × 0.25 mm LD × 0.25 µm) and Agilent automatic injection system

(Agilent GC 7890A and Agilent MS5975C – MSD. The chromatogram was produced by keeping oven temperature at 60 °C for 5 min initially, then the temperature was increased to 90 °C, at a rate of 2 °C/min and then to 90 °C at 5 °C/m for 40 min (Dadalioglu and Evrendilek, 2004). As a result, MSD conditions were as follows: capillary direct interface temperature (280°C), ionization energy (70 eV), mass range (33-330 AMU), EM voltage (Atune + 200) and scan rate (5 scan/s). Herein, helium (He) was used as the gas carrier at a flow rate of 1.0 ml/min with injection of 0.2 µl of oil sample. Quantification was done by an external standard method using calibration curves generated by running GC analysis of representative compounds (Fig. 1). The essential oil components were identified by using the WILEY-NIST MS data library.




Figure 1. Typical gas-chromatogram trace of the essential oil profile extracted from the leaves of the *L. nobilis* L

Concentration–response larvicidal bioassay

Larvicidal bioassays, following the WHO standard protocols instructions (WHO, 2005) for determining the susceptibility or resistance of mosquito larvae to insecticides with suitable modifications, were carried out against the mosquito larvae, treated with the EO extracted from *L. nobilis*. Selected different concentrations of this EO (5, 10, 25 and 35 µl/l) from preliminary toxicological tests, were dissolved in 1 ml acetone and were added separately to the jars, containing 1000 ml of breeding water and 25 larvae of the third or the fourth instar of *Cx. pipiens*. For each larval stage, the experiments were done by using 3 repetitions per concentration with 24 h exposure times (WHO, 2005). Three other jars were left untreated with 1 ml of acetone as negative control replicates. After the exposure time, larvae were removed and placed in clean water. The observed mortality rates were recorded after 24, 48 and 72 h of post-exposure. The observed mortality, in 48 and 72 h, were expressed by the addition of the mortality at 24 h and 48 h, respectively. Dead larvae were identified when they stopped moving after probing the body with a needle.

Statistical analysis

All statistical analyses were performed using Minitab and Prism 7.0 for Windows (Graph Pad Software Inc, USA) with a significance level of $p < 0.05$. The observed percentage mortality was corrected during the assays of the EO larvicidal potentiality. Data have been expressed by the mean \pm standard deviation ($M \pm SD$). All data were analyzed by one-way ANOVA analysis of variance; after angular transformation of observed mortality (%), in order to evaluate the differences between the control and treated series. The mortality (%) of each concentration were analyzed using a Probit model and the lethal concentrations (LC_{50} and LC_{90}), the upper and lower confidential limits and the slope were estimated as described elsewhere (Abbott, 1925; Swaroop et al., 1966).

Reproduction tests

Fecundity experiments were carried out on the eggs of *Cx. pipiens* females emerged from fourth larval stage, survived from treatment with the lethal concentrations estimated after 72 h of delayed mortality ($LC_{50} = 7.18 \mu\text{l/l}$ and the $LC_{90} = 39.08 \mu\text{l/l}$) of the *L. nobilis* EO. For each lethal concentration only 10 females and 10 males were kept in separate breeding cages, and the longevity was recorded until the death of all adults. The laying eggs for each series were collected, counted and transferred to a new jar containing 1000 ml of breeding water for larval hatching. Different parameters of reproduction; the number of laid eggs, hatching rate and the fecundity were studied. The fecundity was calculated by the number of eggs laid in ovitrap divided by number of females let to mate. The significance of *L. nobilis* EO effect on reproduction activity was estimated using Student's t-test.

Results

Yield and chemical composition of *L. nobilis* essential oil

The essential oil extracted by hydrodistillation from leaves of *L. nobilis* exhibited a liquid color change from a dark green to yellow with herbaceous and a woody odor. The result of the obtained leaf-oil yield of the present study was 0.79% (w/w). The chemical composition of the essential oil was analyzed by GC-MS. 39 different components were characterized, during an elution time of 1 h 20 min (Fig. 1), accounting for 99.98% of the total EO composition and these were identified and their quantity were estimated (Table 1). The major EO components were mainly 1,8-cineole (22.41%), Isolongifolene (10.22%), 3-carene (7.74%), Alpha-zingiberene (6.64%), Eremophilene (4.87%), sabinene (4.13%), Euasarone (3.27%), Aristola-1(10), 8-diene (3.09), Gamma-terpinene (2.85%), Delta-selinene (2.36%), 2-carene (2.34%), α -pinene (1.94%), Beta-vatirenene (1.78%) and Beta-caryophyllene (0.36%). Other components existed in traces with less than 1%.

Toxicity assays

The toxicity effects of *L. nobilis* essential oil against the third and fourth instar larvae of *Cx. pipiens* was expressed by the corrected observed mortality, which was recorded at different periods during the treated developmental stage. The lowest used concentration of $5 \mu\text{l/l}$ of the EO caused mortality of 32% for the third stage (L3) and 25% for the fourth stage larvae (L4), after 24 h of exposure time. This observed mortality (%) increased in function of time and doses (Fig. 2). After 72 h, it was noticed that all treated third larvae were dead (100%) with the highest concentration of $35 \mu\text{l/l}$ of the EO, whereas for the fourth ones the mortality for the same dose was 85%. The

data variance analysis showed a significant ($p < 0.001$) insecticidal activity with a concentration-response relationship (Fig. 2). With probit analysis, for the cumulated mortality after 72 h, the lethal concentrations (LC₅₀ and LC₉₀) were estimated with their fiducial limits (95%) from the linear regression curves (Fig. 3) and are listed in Table 2. The non-linear regression curves (Fig. 3), expressed by the mortality probits and the logarithm of *L. nobilis* EO concentrations, reveal a very strong relationship between the corrected mortality (%) and the concentration logarithm for both treated stages. The results of regression analysis of used EO revealed that the mortality rate (Y) is positively correlated with the exposure dose (X) having a regression coefficient (R²) close to 1 in each stage (Table 2). The lethal concentrations were estimated at, LC₅₀ = 3.74 µl/l and the LC₉₀ = 14.47 µl/l for the third larval stage and LC₅₀ was 7.18 µl/l and the LC₉₀ was at 39.08 µl/l, for the fourth larval stage (Table 2).




Figure 2. Recorded mortality (%) with a concentration-response relationship of the treatment with the *L. nobilis* EO applied to newly exuviated larvae of *Cx. pipiens*. (Means \pm SD, $n = 75$) (A = Third instar Larvae, B = fourth instar larvae)




Figure 3. *L. nobilis* essential oil applied to newly exuviated larvae of *Cx. pipiens* at different doses (µl/l): Linear regression dose-response expresses by the probits mortality (%) and the decimal logarithm of *L. nobilis* EO concentrations (µl/l). (A, B = Third and fourth larval stage of *Cx. pipiens* toxicity respectively)

Table 1. Abundance (%) of *L. nobilis* essential oil components identified, using gas chromatography-electron impact mass spectrometry (GC-MS) and their retention time (RT: min)

No.	Compound	Abundance (%)	RT (min)
1	α -pinene	1.943%	8.104
2	sabinene	4.137%	10.199
3	2- β -pinene	1.576%	10.398
4	1,8-cineole	22.417%	14.326
5	b-ocimene	0.482%	14.611
6	γ -terpinene	0.595%	15.217
7	3-carene	7.745%	18.746
8	Gamma.-terpinene	2.857%	25.911
9	2-carene	2.342%	76.703
10	α -caryophyllene	1.289%	76.703
11	α -gurginene	0.697%	76.977
12	Ethyl (e)-cinnamate	1.052%	78.835
13	(e)-germacrene d	1.648%	79.083
14	Aromadendrene	1.630%	79.545
15	Delta.-selinene	2.364%	80.125
16	Alpha-zingiberene	6.694%	80.753
17	Beta-vaterene	0.545%	81.197
18	Alpha-amorphene	1.512%	81.448
19	Gamma-murolene	0.612%	81.551
20	Delta-cadenene	1.149%	81.784
21	Beta-humulene	1.040%	82.468
22	Cis-alpha-bisabolene	2.302%	83.015
23	Euasarone	3.270%	83.953
24	Isolongifolene	10.223%	84.763
25	Neoisolongifolene	0.752%	84.907
26	Bicyclogermacrene	0.752%	85.313
27	Junipene	1.010%	85.742
28	1.1.4.4. Tetrametyl-2-tetralone	0.653%	86.034
29	Valencene 1	0.920%	86.159
30	Caryophyllene	0.815%	86.529
31	Trans-caryophyllene	1.312%	86.669
32	Isoamylcinnamate	0.719%	86.751
33	Ethoxymethyl	1.582%	86.873
34	Caryophylladienol i	0.935%	86.932
35	Beta-vateranene	1.786%	87.238
36	Eremophilene	4.879%	87.582
37	Aristol-1(10),8-diene	3.079%	88.096
38	Dehydroaromadendrene	0.326%	88.469
39	Beta-caryophyllene	0.360%	89.348
Total		99.98%	

Table 2. Toxicity of *L. nobilis* essential oil against Third (L3) and fourth (L4) instar larvae of *Culex pipiens*. Lethal concentrations (LC₅₀ and LC₉₀, Fiducial limits (FL), µl/l)

Larval stage	Regression equation	Slope	LC ₅₀ (µl/l) FL (95%)	LC ₉₀ (µl/l) FL (95%)	R ²	P
L3	y = 2.3899x + 3.3647	0.44	3.33 < 3.74 > 4.33	12.47 < 14.47 > 16.78	0.9009	0.0001
L4	y = 1.772x + 3.4708	0.46	5.93 < 7.18 > 8.68	32.34 < 39.08 > 47.19	0.801	0.0001

Effect of *L. nobilis* essential oil on the longevity of mosquito developmental stages

The result of *L. nobilis* essential EO effect on the longevity of the different developmental stages of *Cx. pipiens* after treatment of the third and fourth instar larvae, with their lethal concentrations (LC₅₀ and LC₉₀) respectively are mentioned in Table 3. The exposure of the new exuviated larvae to the EO caused significant difference (P < 0.05) in the duration of the treated third and fourth larval stage, with both lethal doses compared to control. During the following pupal stage, no difference was recorded in the duration, after treatment with the used lethal concentrations. However, the adult longevity of mosquito species *Cx. pipiens* was considerably reduced (P < 0.005) by the *L. nobilis* EO treatment. The longevity of adults of *Cx. pipiens* emerged from the treated fourth-instar larvae series, was reduced up to 19 days with LC₉₀ and to 21 days for LC₅₀, whereas it was recorded 25 days for the control ones.

Table 3. Effect of the *L. nobilis* EO on the longevity of developmental stages (days); after treatment with the lethal concentrations (LC₅₀ & LC₉₀), of 3rd and 4th instar larvae of *Cx. pipiens*

Stage	Duration of the developmental stages (day) of <i>Cx. pipiens</i> after treatment with lethal concentrations (LC ₅₀ & LC ₉₀).				
	Third instar larvae (L3)			Fourth instar larvae (L4)	
	Control	LC ₅₀ = 3.74 µl/l	LC ₉₀ = 14.47 µl/l	LC ₅₀ = 7.18 µl/l	LC ₉₀ = 39.08 µl/l
L3	4.60 ± 1.33	4.60 ± 2.36	5.16 ± 2.5		
L4	6.15 ± 1.88 ^a	6.80 ± 1.19	7.33 ± 1.13 ^a	7.76 ± 1.33 ^a	8.50 ± 2.66 ^a
Pupae	3.13 ± 1.66	3.43 ± 1.43	3.43 ± 1.33	3.23 ± 1.33	3.23 ± 1.33
Adult	25 ± 1.23 ^{a,b}	21.53 ± 2.33 ^a	20.5 ± 2.23 ^b	21 ± 1.33 ^{a,b}	19.53 ± 3.66 ^{a,b}

The duration (means ± SD) followed by the same letter indicates a significant difference (P < 0.05). (n = 10-70)

Effect of the *L. nobilis* essential oil on reproduction

The effect of *L. nobilis* EO on the female reproduction emerged from the treated fourth instar larvae of *Cx. pipiens*, was evaluated using different parameters (Table 4). Fecundity was highly reduced (P < 0.005) after the treatment with the *L. nobilis* EO. The number of eggs laid was inversely proportional to treatment concentrations. In control females, the number was 672 eggs, whereas it was 583 and 299 eggs, in treated females treated with the two lethal concentrations (LC₅₀ and LC₉₀) respectively (Table 4). The same effect was observed, in the hatching process, with a reduction of the hatching rates. This showed a significant reduction (P < 0.005) according to the increase in the concentration (Table 4). The percentage

of hatching eggs laid by the females emerged from the treated larvae of *Cx. pipiens* was significantly reduced ($p \leq 0.001$) comparatively to control, with a rate of 73.33% with LC₅₀ and 55.2% with CL₉₀, moreover the calculated fecundity percentage showed a decrease at 59.4% and 44.8% for the treated series, with LC₅₀ and LC₉₀ respectively (Table 4).

Table 4. Effect of *L. nobilis* EO on reproduction of the females emerged from the treated fourth instar larvae of *Cx. pipiens* ($n = 10$ females)

Treatment	No. of eggs laid	Hatching rate (%)	Fecundity (%)
Control	672 ^a	98.66 ^a	96.6 ^a
LC ₅₀ = 7.18 µl/l	583 ^{a,b}	73.33 ^{a,b}	59.4 ^{a,b}
LC ₉₀ = 39.08 µl/l	299 ^{a,b}	55.02 ^{a,b}	44.8 ^{a,b}

Numbers followed by the same letter indicate a significant difference ($P < 0.005$)

Discussion

Yield and chemical composition of the essential oil

The essential oil extraction from dry *L. nobilis* leaves plant, collected from the study area (high-plains region, Setif, Algeria), demonstrated that the essential oil yield was 0.79% (w/w) which confirm the result previously reported (Elharas et al., 2013). Whereas the EO yield of the same plant from Tebessa (Northeast Algeria) was 0.96% (Bouzidi et al., 2020). These EO yields from the Algerian local plants of *L. nobilis*, were higher than the yield extracted from *L. nobilis* from Montecorice, Italy (0.57%) (Caputo et al., 2017), Tunisia (0.58%) and Morocco (0.65%) (Mediouni-BenJemaâ et al., 2012). While this finding showed a lower yield than those reported elsewhere, with the value of 1.13% (de Oliveira et al., 2021) and 1.3% (Ouibrahim et al., 2013). The varied essential oil yield is based on the quality of the used plant material, combined with other factors, such as the growth stage, soil quality, climate conditions, time of harvest and drying period (Mohammedi et al., 2020). Indeed, methods of drying aromatic and medicinal plants can affect the yield and the chemical composition percentages, mainly in the quantitative aspect of essential oils (Sellami et al., 2011; Abdelmajeed et al., 2013). The non-dried plant under good conditions may be degraded, and consequently can lose some of its entire essential oils (Sarić-Krsmanović et al., 2020). As such, reports on the chemical compositions of *L. nobilis* EOs indicated variations both quantitatively and qualitatively. This study, for instance, identified 39 EO components, or 99.98% of the total oil content, whereas a previous study (Caputo et al., 2020) has identified 55 compounds accounting for 91.6% of the total essential oil; including 1.8 cineole (31.9%), sabinene (12.2%), linalool (10.2%), α -terpinylacetate (5.9%), α -pinene (5.8%), α -terpineol (3.3%), methyl-eugenol (3.3%), neoiso-isopulegol (2.5%), eugenol (1.6%), β -pinene (1.4%) and γ -terpinene (1.0%). This mixture of chemicals is almost similar qualitatively to the one reported previously (Marzouki et al., 2009) but still incompatible with our sample.

Based on the recent available literature, the comparison of the chemical composition of *L. nobilis* essential oil under investigation and that of other areas showed substantial similarities and differences. The percentage of 1,8-cineole was found as 22.41%, which is closely related to EOs observed in the laurel plants from Algeria (24.65%) (Miliani et al., 2017), Greek (30.08%) and Gorgia (29.2%) (Galina et al., 2020). Moreover, higher

percentages of 1.8 cineole were recorded from some Mediterranean countries, Turkey (44.97%), Tunisia (56.0%) (Nabila et al., 2020) and Morocco (52.43%) (Nafis et al., 2020). In the present work the amount of other constituents, such as the sabinene was found 4.137% which is higher than that found in the essential oils yield previously reported and lower than that reported elsewhere (Miliani et al., 2017) with 6.13%. Also, it is important to notice that the linalool component that was found with different percentages in previous studies (Miliani et al., 2017), was absent in our analysis. This result is in accordance with other analysis where linalool has not been found (Derwich et al., 2009). The content of 1.8-cineole (22.41%) is by far the most dominant component of *L. nobilis* EO from Setif area (Northeast Algeria) besides that of Tizi Ouzou (Northeast Algeria) exhibiting 35.5% of the total essential oil composition (Mohammedi et al., 2020). Beside the difference of the plants geographical origins (Sangun et al., 2007), the reported differences in the EO composition and yield, in this study, might be attributed to environmental conditions, harvest and postharvest processing factors (Galina et al., 2020).

Insecticidal activity of L. nobilis L. essential oil

Use of natural plant origin products, with insecticidal properties have been encouraged in the control of a variety of insect pests and vectors. Mosquitoes in the larval stage are the main attractive targets for control because they breed in water, and thus, it is easy to deal with them in this habitat. Many studies have reported the plant extract larvicidal effect against mosquito larvae (Prabhu et al., 2011). The crude extract obtained from the *Leucas aspera* leaf showed a larvicidal activity against *Culex quinquefasciatus*, *Aedes aegypti* larvae (Maheswaran et al., 2008; Govindarajan, 2009) and *Anopheles stephensi* larvae (Govindarajan, 2011). Other works have reported the *Cassia fistula* extracts of the larvicidal toxicity and ovicidal and repellent activities against *Aedes aegypti* (Govindarajan, 2009). The *Acalypha indica* and *Citrullus vulgaris* leaf extract effectiveness was tested too on larvicidal, ovicidal activity against *Anopheles stephensi* (Mullai et al., 2008). The bioassays using the *L. nobilis* essential oil, against the larval stages of the mosquito species *Cx. pipiens* showed a larvicidal activity expressed by a high larval mortality of the treated series compared to the control ones. The difference in sensitivity to plant extracts appears within the mosquito genus (genera: *Culex*, *Aedes*, *Ochlerotatus*, or *Anopheles*) that would be caused by behavioral and physiological variations of different species (Saxena and Tikku, 1990). The *L. nobilis* EO exhibits also high effective larvicidal activity in the newly exuviated third and fourth stage larvae of *Cx. pipiens*, along with mortality rates range with a dose-response relationship. the same efficacy was reported using the same plant extract against *Culiseta longiareolata* (Elharas et al., 2013).

In the present study, the *L. nobilis* essential oil treatment reduced the larval duration, of the third and fourth larval stages. Therefore the developmental time of pupal and the emerged adult were not affected. Also the emerged adults, showed a significant reduction in their longevity. The adult which emerged from treated larvae were morphologically normal but showed a great reduction in fecundity. The same results were mentioned against *Aedes aegypti*, using EO plants (Warikoo, et al., 2011) and *Cx. pipiens* (Djeghader, et al., 2018; Djeddar, et al., 2021) using plant extracts. From the present study it was concluded that the *L. nobilis* EO showed a promising larvicidal agent against mosquito larvae and also reduced the longevity of different developmental stages, egg productions and fecundity.

Conclusion

The essential oil extracted from *L. nobilis* with an interesting yield contains many important components. The bioassay investigation confirm that this EO exhibited potent larvicidal activity, a development disturbance and reduction in fecundity against *Cx. pipiens*. This plant essential oil proves to be a promising alternative as an efficient bio-insecticidal agent. The potential effect of this essential oil is reasonable with the presence of beneficial phytochemicals like 1.8-cineole, isolongifolene, 3-carene and sabinene as well as some minor constituents, which may play a disruptive role.

Acknowledgments. The authors would like to thank the laboratory team of Molecular Biology and Genetics Department of Istanbul University, Turkey, for their help for the CPG-MS analyses and MERS, Algeria, for the financial support.

REFERENCES

- [1] Abbott, W. S. (1925): A method of computing the effectiveness of an insecticide. – *Journal of Economic Entomology* 18: 265-267.
- [2] Abdelmajeed, N. A., Danial, E. N., Ayad, H. S. (2013): The effect of environmental stress on qualitative and quantitative essential oil of aromatic and medicinal plants. – *Archives des Sciences* 66: 100-120.
- [3] Aïssaoui, L., Boudjelida, H. (2017): Diversity and distribution of culicinae fauna in Tebessa district (North-East of Algeria). – *International Journal of Mosquito Research* 4: 7-12.
- [4] Amalraj, D. D., Sahu, S. S., Jambulingam, P., Doss, P. S. B., Kalyanasundaram, M., Das, P. K. (2000): Efficacy of aqueous suspension and granular formulations of *Bacillus thuringiensis* (Vectobac) against mosquito vectors. – *Acta Tropica* 75: 243-246.
- [5] Arroussi, D. E. R., Bouaziz, A., Boudjelida, H. (2021): Mosquito survey reveals the first record of *Aedes* (Diptera: Culicidae) species in urban area, Annaba district, Northeastern Algeria. – *Polish Journal of Entomology* 90(1): 14-26. DOI: 5604/01.3001.0014.8065.
- [6] Bouaziz, A., Amira, K., Djeghader, N., Aïssaoui, L., Boudjelida, H. (2017): Impact of an insect growth regulator on the development and the reproduction potency of mosquito. – *Journal of Entomology and Zoology Studies* 5: 1662-1667.
- [7] Boudjelida, H., Bouaziz, A., Thomas, S., Smagghe, G., Soltani, N. (2005): Effects of ecdysone agonist halofenozide against *Culex pipiens*. – *Pesticide Biochemistry and Physiology* 83: 115-123.
- [8] Bouzidi, O., Tine, S., Hamaidia, K., Tine-Djebbar, F., Soltani, N. (2020): Chemical Composition and Bioefficacy of Essential Oil from Bay Laurel Shrub (Laurales: Lauraceae) against *Culiseta longiareolata* (Macquart) (Diptera: Culicidae) Larvae. – *Journal of Entomological Science* 55(2): 262-272.
- [9] Caputo, L., Nazzaro, F., Souza, L. F., Aliberti, L., De Martino, L., Fratianni, F., Coppola, R., De Feo, V. (2017): *Laurus nobilis*: composition of essential oil and its biological activities. – *Molecules* 22: 930-938.
- [10] Caputo, L., Smeriglio, A., Trombetta, D., Cornara, L., Trevena, G., Valussi, M., Fratianni, F., De Feo, V., Nazzaro, F. (2020): Chemical composition and biological activities of the essential oils of *Leptospermum petersonii* and *Eucalyptus gunnii*. – *Front Microbiology* 11: 409-416.
- [11] Corzo, F. L., Traverso, L., Sterkel, M., Benavente, A., Ajmat, M. T., Ons, S. (2020): *Plodia interpunctella* (Lepidoptera: Pyralidae): Intoxication with essential oils isolated from *Lippia turbinata* (Griseb.) and analysis of neuropeptides and neuropeptide receptors,

- putative targets for pest control. – Archives of Insect Biochemistry and Physiology 104: 21684-21692.
- [12] da Costa, G. M., dos Santos, M. C. C., Everton, G. O., de Sousa, P. de C. D., Rangel, M. E. S. (2020): Physicochemical properties, toxicity and larvicidal activity of the essential oil of *Cymbopogon winterianus* in front of *Aedes aegypti*. – Ciência Nature 42: 39-47.
- [13] Dadalıođlu, I., Evrendilek, G. A. (2004): Chemical compositions and antibacterial effects of essential oils of Turkish oregano (*Origanum minutiflorum*), bay laurel (*Laurus nobilis*), Spanish lavender (*Lavandula stoechas* L.), and fennel (*Foeniculum vulgare*) on common foodborne pathogens. – Journal of Agricultural and Food Chemistry 52: 8255-8260.
- [14] Dar, A. A., Jamal, K. (2021): Evaluation of larvicidal effects of aqueous leaf and flower extracts of *Ammi majus* (Linn.) on mosquito larvae. – Munis Entomology and Zoology 16(2): 891-897.
- [15] de Oliveira, M. S., da Silva, V. M. P., Freitas, L., Silva, S. G., Cruz, J., Andrade, E. H. de A. (2021): Extraction yield, chemical composition, preliminary toxicity of *Bignonia nocturna* (Barb. Rodr.) LG Lohmann (Bignoniaceae) essential oil and in silico evaluation of the interaction. – Chemistry and Biodiversity 18(4): e2000982.
- [16] Derwich, E., Benziane, Z., Boukir, A. (2009): Chemical composition and antibacterial activity of leaves essential oil of *Laurus nobilis* from Morocco. – Australian Journal of Basic and Applied Sciences 3: 3818-3824.
- [17] Djeddar, H., Boudjelida, H., Arroussi, D. E. R. (2021): New alternative for culicidian fauna control using *Borago officinalis* and *Drimia maritima* plant extracts. – Biodiversitas 22(12): 5688-5694.
- [18] Djeghader, N. E., Aïssaoui, L., Amira, K., Boudjelida, H. (2018): Toxicity evaluation and effects on the development of a plant extract, the Saponin, on the domestic mosquito, *Culex pipiens*. – International Journal of Mosquito Research 5: 1-5.
- [19] Dmitry, A. K., Naida, M. A. (2019): Phenolic compounds of *Laurus nobilis* (review). – Pharmacy and Pharmacology 7(5): 244-259. DOI: 10.19163/2307-9266-2019-7-5-244-259.
- [20] Elharas, K., Daagare, A., Mesifioui, A., Ouhssine, M. (2013): Antibacterial activity of essential oil of inflorescence *laurus nobilis* and *lavandula angustifolia*. – Afric Science 9: 134-141.
- [21] Flores, H. A., O'Neill, S. L. (2018): Controlling vector-borne diseases by releasing modified mosquitoes. – Nature Reviews Microbiology 16: 508-518.
- [22] Galina, S., Tanya, G., Velizar, G., Magdalena, S., Zhana, P., Albena, S., Valcho, D. Z. (2020): Comparative study on the chemical composition Greece and Gorgia and the antibacterial activity of their essential oil. – Heliyon 6(12): 1-6.
- [23] Georgiev, E., Stoyanova, A. (2006): A Guide for the Specialist in the Aromatic Industry. – UFT Academic Publishing House, Plovdiv.
- [24] Govindarajan, M. (2009): Bioefficacy of *Cassia fistula* Linn. (Leguminosae) leaf extract against chikungunya vector, *Aedes aegypti* (Diptera: Culicidae). – European Review for Medical and Pharmacological Sciences 13(2): 99-103.
- [25] Govindarajan, M. (2011): Evaluation of indigenous plant extracts against the malarial vector, *Anopheles stephensi* (Liston) (Diptera: Culicidae). – Parasitology Research.109: 93-103.
- [26] Govindarajan, M., Sivakumar, R., Rajeswari, M., Yogalakshmi, K. (2012): Chemical composition and larvicidal activity of essential oil from *Mentha spicata* (Linn.) against three mosquito species. – Parasitology Research 110: 2023-2032.
- [27] Gunasekaran, K., Doss, P. S. B., Vaidyanathan, K. (2004): Laboratory and field evaluation of Teknar HP-D, a biolarvicidal formulation of *Bacillus thuringiensis* sp. israelensis, against mosquito vectors. – Acta Tropica 92: 109-118.

- [28] Kilic, A., Hafizoglu, H., Kollmannsberger, H., Nitz, S. (2004): Volatile constituents and key odorants in leaves, buds, flowers, and fruit of *Laurus nobilis* L. – Journal of Agricultural and Food Chemistry 52: 1601-1606.
- [29] Lounibos, L. P. (2002): Invasions by insect vectors of human disease. – Annual Review of Entomology 47: 233-266.
- [30] Maheswaran, R., Sathish, S., Ignacimuthu, S. (2008): Larvicidal activity of *Leucas aspera* (Wild.) against the larvae of *Culex quinquefasciatus* Say. and *Aedes aegypti* L. – International Journal of Integrative Biology 2008(2): 214-217.
- [31] Marzouki, H., Piras, A., Salah, K. B. H., Medini, H., Pivetta, T., Bouzid, S., Falconieri, D. (2009): Essential oil composition and variability of *Laurus nobilis* L. growing in Tunisia, comparison and chemometric investigation of different plant organs. – Natural products Research 23(4): 343-54.
- [32] Marzouki, H., Khaldi, A., Marongiu, B., Piras, A., Harzallah-Skhiri, F. (2011): Chemical polymorphism of essential oils from populations of *Laurus nobilis* grown on Tunisia, Algeria and France. – Natural Product Communications 6(10): 1483-1486.
- [33] Mediouni-BenJemaâ, J., Tersim, N., Taleb-Toudert, K., Larbi-Khouja, M. (2012): Insecticidal activities of essential oils from leaves of *Laurus nobilis* L. from Tunisia, Algeria and Morocco, and comparative chemical composition. – Journal of Stored Products Research 48: 97-104.
- [34] Miliani, A., Boukhatem, M., Saidi, F. (2017): Chemical composition and antimicrobial activity of the Algerian *Laurus nobilis* essential oil. – Algerian Journal Natural Products 5(2): 507-514.
- [35] Minetti, C., Ingham, V. A., Ranson, H. (2020): Effects of insecticide resistance and exposure on Plasmodium development in *Anopheles* mosquitoes. – Current Opinion of Insect Sciences 39: 42-49.
- [36] Mohammedi, H., Mecherara-Idjeri, S., Hassani, A. (2020): Variability in essential oil composition, antioxidant and antimicrobial activities of *Ruta montana* L. collected from different geographical regions in Algeria. – Journal of Essential Oil Research 32: 88-101.
- [37] Mouhamadou, C. S., Luan, K., Fodjo, B. K., West, A. J., McCord, M. G., Apperson, C. S., Roe, R. M. (2020): Development of an insecticide-free trapping bednet to control mosquitoes and manage resistance in malaria vector control: a new way of thinking. – Insects: 11(11): 732-753.
- [38] Mullai, K., Jebanesan, A., Pushpanathan, T. (2008): Effect of bioactive fractions of *Citrullus vulgaris* Schrad. leaf extract against *Anopheles stephensi* and *Aedes aegypti*. – Parasitology Research 102(5): 951-955.
- [39] Muturi, E. J., Ramirez, J. L., Doll, K. M., Bowman, M. J. (2017): Combined toxicity of three essential oils against *Aedes aegypti* (Diptera: Culicidae) larvae. – Journal of Medical Entomology 54: 1684-1691.
- [40] Nabila, B., Piras, A., Fouzia, B., Falconieri, D., Kheira, G., Fedoul, F. F., Majda, S. R. (2020): Chemical composition and antibacterial activity of the essential oil of *Laurus nobilis* leaves. – Natural Product Research 1: 1-5.
- [41] Nabti, I., Bounechada, M. (2019): Larvicidal activities of essential oils extracted from five algerian medicinal plants against *Culiseta longiareolata* Macquart. larvae (Diptera: Culicidae). – European Journal of Biology 78: 133-138.
- [42] Nafis, A., Kasrati, A., Jamali, C. A., Custódio, L., Vitalini, S., Iriti, M., Hassani, L. (2020): A comparative study of the in vitro antimicrobial and synergistic effect of essential oils from *Laurus nobilis* L. and *Prunus armeniaca* L. from Morocco with antimicrobial drugs: new approach for health promoting products. – Antibiotics 9: 140-148.
- [43] Ouibrahim, A., Tlili-Ait-kaki, Y., Bennadja, S., Amrouni, S., Djahoudi, A. G., Djebbar, M. R. (2013): Evaluation of antibacterial activity of *Laurus nobilis* L., *Rosmarinus officinalis* L. and *Ocimum basilicum* L. from Northeast of Algeria. – African Journal of Microbiological Research 7: 4968-4973.

- [44] Patrakar, R., Mansuriya, M., Patil, P. (2012): Phytochemical and pharmacological review on *Laurus nobilis*. – International Journal of Chemistry and Pharmaceutical Sciences 1: 595-602.
- [45] Prabhu, K., Murugan, K., Nareshkumar, A., Ramasubramanian, N., Bragadeeswaran, S. (2011): Larvicidal and repellent potential of *Moringa oleifera* against malarial vector, *Anopheles stephensi* Liston (Insecta: Diptera: Culicidae). – Asian Pacific Journal of Tropical Biomedicine 2011(1): 127-132.
- [46] Raja, T. R. W., Ganesan, P., Gandhi, M. R., Duraipandiyan, V., Paulraj, M. G., Balakrishna, K., Al-Dhabi, N. A., Ignacimuthu, S. (2018): Effect of compound Musizin isolated from *Rhamnus wightii* Wight and Arn on the immature stages of filarial vector mosquito *Culex quinquefasciatus* Say (Diptera: Culicidae) and its non-target studies. – Biocatalysis and Agricultural Biotechnology 16: 37-42.
- [47] Rawani, A., Ghosh, A., Chandra, G. (2010): Mosquito larvicidal activities of *Solanum nigrum* L. leaf extract against *Culex quinquefasciatus* Say. – Parasitology Research 107: 1235-1240. DOI 10.1007/s00436-010-1993-9).
- [48] Richards, S. L., Byrd, B. D., Reiskind, M. H., White, A. V. (2020): Assessing insecticide resistance in adult mosquitoes: perspectives on current methods. – Environmental Health Insights 14: 1-7.
- [49] Salavati, B., Zahirnia, A. H., Nasirian, H., Azari-Hamidian, S. (2021): Trend of mosquito (Diptera: Culicidae) monthly distribution in Sanandaj County of Iran. – Biodiversitas 22(11): 4705-4715.
- [50] Sangun, M. K., Aydin, E., Timur, M., Karadeniz, H., Caliskan, M., Ozkan, A. (2007): Comparison of chemical composition of the essential oil of *Laurus nobilis* L. leaves and fruits from different regions of Hatay, Turkey. – Journal Environmental Biology 28: 731-733.
- [51] Sarić-Krsmanović, M., Dragumilo, A., Gajić Umiljendić, J., Radivojević, L., Šantrić, L., Đurović-Pejčev, R. (2020): Infestation of field dodder (*Cuscuta campestris* Yunck.) promotes changes in host dry weight and essential oil production in two aromatic plants, peppermint and chamomile. – Plants 9: 1286-1295.
- [52] Saxena, B. P., Tikku, K. (1990): Impact of natural products on the physiology of phytophagous insects. – Proceedings: Animal Science 99: 185-198.
- [53] Sellami, I. H., Wannas, W. A., Bettaieb, I., Berrima, S., Chahed, T., Marzouk, B., Limam, F. (2011): Qualitative and quantitative changes in the essential oil of *Laurus nobilis* L. leaves as affected by different drying methods. – Food Chemistry 126: 691-697.
- [54] Smith, C. J., Perfetti, T. A. (2020): A comparison of the persistence, toxicity and exposure to high-volume natural plant-derived and synthetic pesticides. – Toxicology Research and Application 4: 1-15.
- [55] Swaroop, S., Gilroy, A. B., Uemura, K., Organization, W. H. (1966): Statistical Methods in Malaria Eradication. – World Health Organization, Geneva.
- [56] Warikoo, R., Wahab, N., Kumar, S. (2011): Oviposition-altering and ovicidal potentials of five essential oils against female adults of the dengue vector, *Aedes aegypti* L. – Parasitology Research 109(4): 1125-1131.
- [57] WHO (World Health Organization). (1991): Insect and Rodent Control through Environmental Management: A Community Action Program. – WHO, Geneva, pp. 29-34.
- [58] WHO (World Health Organization). (2005): Guidelines for Laboratory and Field Testing of Mosquito Larvicides. – World Health Organization, Geneva.
- [59] Wilson, A. L., Courtenay, O., Kelly-Hope, L. A., Scott, T. W., Takken, W., Torr, S. J., Lindsay, S. W. (2020): The importance of vector control for the control and elimination of vector-borne diseases. – PLoS Neglected Tropical Diseases 14(1): 7831-7838.